



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2772106 C 2017/11/07

(11)(21) **2 772 106**

(12) **BREVET CANADIEN  
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2010/08/30  
(87) Date publication PCT/PCT Publication Date: 2011/03/03  
(45) Date de délivrance/Issue Date: 2017/11/07  
(85) Entrée phase nationale/National Entry: 2012/02/24  
(86) N° demande PCT/PCT Application No.: CA 2010/001253  
(87) N° publication PCT/PCT Publication No.: 2011/022809  
(30) Priorité/Priority: 2009/08/28 (US61/237,906)

(51) Cl.Int./Int.Cl. *C12N 15/31* (2006.01),  
*A01N 63/04* (2006.01), *A01P 3/00* (2006.01),  
*C07K 14/37* (2006.01), *C12N 1/14* (2006.01),  
*C12N 15/63* (2006.01), *C12N 15/80* (2006.01),  
*C12Q 1/68* (2006.01)


(72) Inventeur/Inventor:  
VUJANOVIC, VLADIMIR, CA

(73) Propriétaire/Owner:  
UNIVERSITY OF SASKATCHEWAN, CA

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : LUTTE BIOLOGIQUE CONTRE FUSARIUM ET LES MYCOTOXINES DE FUSARIUM

(54) Title: FUSARIUM AND FUSARIUM MYCOTOXIN BIOCONTROL



(57) Abrégé/Abstract:

The present invention relates to a novel ascomyceteous fungus, *Sphaerodes mycoparasitia* strain IDAC 301008-01, for controlling *Fusarium* plant pathogens, disease symptoms, and mycotoxins in planta and ex planta. Uses, methods, compositions, sequences, and products are also disclosed herein.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPO 191

OPIC



CIPO

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
3 March 2011 (03.03.2011)

PCT

(10) International Publication Number  
**WO 2011/022809 A1**

## (51) International Patent Classification:

*C12N 15/31* (2006.01)    *C12N 1/14* (2006.01)  
*A01N 63/04* (2006.01)    *C12N 15/63* (2006.01)  
*A01P 3/00* (2006.01)    *C12N 15/80* (2006.01)  
*C07K 14/37* (2006.01)    *C12Q 1/68* (2006.01)

## (21) International Application Number:

PCT/CA2010/001253

## (22) International Filing Date:

30 August 2010 (30.08.2010)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

61/237,906    28 August 2009 (28.08.2009)    US

(71) Applicant (for all designated States except US): **UNIVERSITY OF SASKATCHEWAN** [CA/CA]; Office of Vice-President of Research, Industry Liaison Office, Box 5000 RPO University, Saskatoon, Saskatchewan S7N 1 K2 (CA).

## (72) Inventor; and

(75) Inventor/Applicant (for US only): **VUJANOVIC, Vladimir** [CA/CA]; 211-301 Tait Place, Saskatoon, Saskatchewan S7H 5L5 (CA).

(74) Agents: **POLONENKO, DANIEL** et al.; Gowling Lafleur Henderson LLP, Box 30, 2300-550 Burrard Street, Vancouver, British Columbia V6C 2B5 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,

[Continued on next page]

(54) Title: FUSARIUM AND FUSARIUM MYCOTOXIN BIOCONTROL




Figure 1

(57) Abstract: The present invention relates to a novel ascomyceteous fungus, *Sphaerodes mycoparasitia* strain IDAC 301008-01, for controlling *Fusarium* plant pathogens, disease symptoms, and mycotoxins *in planta* and *ex planta*. Uses, methods, compositions, sequences, and products are also disclosed herein.

**WO 2011/022809 A1** 

---

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG). **Published:**  
— *with international search report (Art. 21(3))*  
— *with sequence listing part of description (Rule 5.2(a))*

## **FUSARIUM AND FUSARIUM MYCOTOXIN BIOCONTROL**

### FIELD OF THE INVENTION

The present invention relates to novel biocontrol agents, related compositions with mycocidal effect and uses thereof. In particular, the present agent has a mycocidal effect on fungi of the genus *Fusarium*. The present invention further relates to the isolated fungal inoculant, genes, proteins and/or organisms as well as uses, methods, compositions, involving the same.

### BACKGROUND TO THE INVENTION

*Fusarium* is a filamentous fungus widely distributed on plants and in the soil. Certain *Fusarium* species are plant pathogens. For example, *Fusarium oxysporum*, causes *Fusarium* wilt disease in more than a hundred species of plants. It does so by colonizing the plant xylem which can result in blockage and breakdown. When this occurs symptoms such as leaf wilting and yellowing appear in the plant eventually leading to the plant's death. This condition was the primary cause of the decline and disappearance of the Gros Michel banana cultivar from markets around the world. Recently a new strain has begun attacking plants of the dominant Cavendish cultivar leading to fears that, in the absence of a solution, this cultivar will too disappear from world markets.

*Fusarium* root rot is a major cause of seedling mortality in forest nurseries and also causes reduced survival after outplanting during the first growing season. The disease is caused by several *Fusarium* species and is common in many parts of the world. The disease is a particular problem in Western Canada and the United States and also in the North Central and Southern States. In addition, Pine pitch canker - caused by the fungus *Fusarium circinatum* - is a serious disease of pine trees and a threat to the forest industry in particularly in the US and New Zealand. Radiata or Monterey pine is highly susceptible to the disease with mortality rates in mature trees reaching 80% in some areas of California.

*Fusarium* Head Blight (FHB), also known as 'ear blight' or 'scab', is a disease of wheat, barley, oats and other small cereal grains caused by *Fusarium*. Corn and maize can be affected

by a similar condition known as 'ear rot'. The aforementioned condition can reduce the yield and grade of the crop, and can potentially contaminate the grain with mycotoxins. It is estimated that FHB costs the cereal industry almost \$5 billion annually. In recent years, FHB has proved a significant and growing problem for the commercially important wheat and barley crops in Western Canada. In the past *Fusarium oxysporum* and *Fusarium avenaceum* have been identified as the two species primarily associated with FHB in Canada wherein yields in some affected fields have been reduced by 30% or more. FHB and ear rot harvested grain is often contaminated with mycotoxins such as deoxinivalenol (DON) trichothecenes associated with feed refusal, general digestive disorders, diarrhea, hemorrhages. The emergence of the toxigenic 3-acetyldeoxynivalenol (3-ADON) *Fusarium graminearum* population in North America is a fairly recent phenomenon. It is replacing 15-ADON chemotype. Moreover, the mycotoxin profiles of *F. culmorum* - similar to *F. graminearum* under both laboratory and field conditions-represents an additional threat for corn, maize and wheat production worldwide. Currently, there are no effective control measures for FHB and associated mycotoxins. Additionally, there are also no resistant varieties of corn, wheat, barley, oats, or other small grain cereals. Fungicides can be effective but only temporarily suppress the disease.

In addition to being a common plant pathogen, *Fusarium* spp. can opportunistically infect animals and can be the causative agents of superficial and/or systemic infections in humans. *Fusaria* are one of the most drug-resistant fungi making *Fusaria* infections difficult to treat and invasive infections can prove fatal.

#### SUMMARY OF THE INVENTION

The present invention relates to a novel ascomyceteous fungus, *Sphaerodes mycoparasitia* identified as strain IDAC 301008-01 and alternatively as strain SMCD2220-01, that has a mycoparasitic and antimycotoxin effect. In addition, mycotoxic proteins have been isolated from *S. mycoparasitia* strain IDAC 301008-01. Uses, methods, compositions, sequences, and products are also disclosed herein.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a micrograph showing *Sphaerodes mycoparasitica* cultures after two weeks of incubation on (A) Modified Leonian's agar and (B) Potato dextrose agar—upper sides, (C) and (D) - down sides;

Figure 2 is a micrograph showing *Sphaerodes mycoparasitica* ascospore germination, showing single-polar and double-polar germination patterns as well as hyphal anastomosis (arrow) formation pattern;

Figure 3 is a micrograph showing *Sphaerodes mycoparasitica*: (A) Ascoma, (B) Hyaline seta arising from the neck (arrow), (C) Reticulate ascospores (arrows), (D) Smooth ascospore (arrow), (E) Triangular ascospore (arrow), (F) Phialides produced on ascoma surrounding hyphae, (G) Ampulliform phialide arising from the surface of ascoma peridial wall, (H) Formation of mature and starting ascomata, (I) Formation of hook-like structures by *S. mycoparasitica* parasitising on living hypha of *Fusarium oxysporum* (arrows) and (J) Large view of hook-like structure on living hypha of *F. avenaceum*. Bar scales for (A) and (H) are 50 pm; for (B), (C), (D), (E), (F), (G) and (J) are 10 pm; for (I) 25 pm;

Figure 4 (A) is a micrograph showing *Sphaerodes mycoparasitica* ascospores showing the conspicuous wall ornamentation and prominent irregular longitudinal ribs (arrows); (B) shows *Sphaerodes quadrangularis* ascospores. Bar scales 5 pm;

Figure 5(a) is a micrograph showing inhibition of mycelial growth of (A) *Fusarium oxysporum* and (B) *Fusarium graminearum* by extracellular proteins recovered from a *Sphaerodes mycoparasitica* culture; 5(b) is a chart showing the percent inhibition of mycelial growth of A) *Fusarium oxysporum* and (B) *Fusarium graminearum* by *S. mycoparasitica* extracellular proteins;

Figure 6 is a FPLC chromatogram showing the f1 and f2 extracellular proteins recovered from a *Sphaerodes mycoparasitica* culture;

Figure 7 is micrograph of SDS-PAGE gels of purified extracellular proteins recovered from a *Sphaerodes mycoparasitica* culture. Lane M contains marker proteins. Lane 1 contains the purified protein from peak f1. Lane 2 contains the purified protein from peak f2;

Figure 8 are micrographs showing the inhibition of *F. oxysporum* spore germination (A-a-b) and *F. graminearum* spore germination (B-1-b) by purified proteins f1 and f2 compared to controls (A-c and B-c);

Figure 9 is a chart showing the level of 3-ADON degradation by *Sphaerodes mycoparasitica* in potato-dextrose broth analysed by HPLC;

Figure 10 is a schematic illustration of a system for growing wheat plants in a container (4 x 4 x 16 cm) with different layers of soil-less growing mixes;

Figure 11 is a micrograph of a gel showing SmyITSF/R primers amplified PCR products for *S. mycoparasitica* (SM), five *Fusarium* strains (Fa = *F. avenaceum*, Fo = *F. oxysporum*, Fs = *F. sporotrichioides*, Fg3 = *F. graminearum* chemotype 3, and Fg15 = *F. graminearum* chemotype 15), two *Trichoderma* species (T22 = *T. harzianum* T22 and Tv = *T. viride*), two *Cladosporium* species ACC = *C. cladosporioides* and CM = *C. minourae*), and *Penicillium aurantiogriseurn* (PA) were electrophoresed on 1% agarose gel at 100 V for 20 minutes. The size of the band is around 300 to 400 bp;

Figure 12 are charts showing standard linear curves for (A) *Sphaerodes mycoparasitica* (in the range of  $3.8 \times 10^2$  to  $3.8 \times 10^{-2}$  ng in ten-fold decreasing manner); (B) *Fusarium graminearum* 3-ADCIN (in the range of  $2.7 \times 10^3$  to  $2.7 \times 10^{-1}$  ng in ten-fold decreasing manner); (C) *Trichoderma harzianum* T-22 (in the range of  $7.0 \times 10^2$  to  $7.0 \times 10^{-2}$  ng in ten-fold increasing manner);

Figure 13 is a chart showing RT-PCR sigmoidal coloured curves for *Sphaerodes mycoparasitica* (SMCD 2220-01), with 0.025 fluorescence line, in the ranges of  $3.8 \times 10^2$  to  $3.8 \times 10^{-2}$  ng in a ten-fold decreasing manner;

Figure 14 are charts showing quantities of genomic DNAs for (A) *F. graminearum* (Fgra); (b) *S. mycoparasitica* (SM); and (C) *Trichoderma harzianum* (T-22) monitored in spring

wheat roots using genus-specific quantification real-time PCR. All values were means of 6 replicates. Error bars indicate SD;

Figure 15 is a chart showing real-time fluorescence curves of *tri5* gene sequences amplified by using Tox5- 1/2 primer set from total DNA extracted from dual-culture assays of *Fusarium graminearum* strains and pre-inoculated *Sphaerodes mycoparasitica* SMCD2220-01 (SNI) or singly grown *F. graminearum* 3-ADON and 15-ADON chemotypes;

Figure 16 are micrographs of *Sphaerodes mycoparasitica* ascomata and ascospores: (A) formation of *S. mycoparasitica* ascomata on a colony of *F. avenaceum* (Arrows indicate that ascomata were produced near or surrounding the *Fusarium* culture); (B) production of numerous *S. mycoparasitica* ostiolated perithecia being produced on a *F. oxysporum* colony (Arrows indicate perithecia were formed on the *Fusarium* isolate); (C) ungerminated dark-brownish reticulated *S. mycoparasitica* ascospore; (D) germinating ascospore of *S. mycoparasitica* in *F. oxysporum*-filtrate suspension showing one polar germ pore; (E) single polar germinating spore of *S. mycoparasitica* after 3d suspension in a *F. oxysporum*-filtrate; (F) ascospore of *S. mycoparasitica* illustrating two polar germination in *F. oxysporum*-filtrate suspension after 3d with an additional 1d on PDA; (G) pattern of *S. mycoparasitica* two polar germination in *F. avenaceum*-filtrate suspension with additional 1d on PDA; (H) single polar germination in *S. mycoparasitica* spore suspended 3d in *F. avenaceum*-filtrate with 1d incubation on PDA; (I) single and double polar germinations demonstrated by *S. mycoparasitica* ascospores after 3d incubation in a *F. avenaceum*-filtrate suspension plus 1 additional day incubation on PDA. Scale bars for (C) to (H) are 10µm and for (I) is 20µm. SM, Fa, and Fo represent *S. mycoparasitica*, *F. avenaceum*, and *F. oxysporum*, respectively;

Figure 17 is a chart showing spore germination patterns of *Sphaerodes mycoparasitica* biotrophic mycoparasitic fungus with spores isolated from a *F. oxysporum* colony. Single (■) and two (□) polar germination in a *F. avenaceum*-filtrate suspension; and Single (▣) and two (■) polar germination in a *F. oxysporum*-filtrate suspension. 1 = 1d suspension, 2 = 1d suspension plus 1d on PDA incubation, 3 = 3d suspension, and 4 = 3d suspension plus 1d on PDA incubation. Each incubation day for *S. mycoparasitica* was analyzed separately;



Figure 18 is a chart showing germination of *Sphaerodes mycoparasitica* ascospores in filtrates of six *Fusarium* strains and water suspension treatments at four different incubation days. Suspension in different treatments were: Fave-filtrate = *F. avenaceum*-filtrate; Foxy-filtrate = *F. oxysporum*-filtrate; Fgra3-filtrate = *F. graminearum* chemotype 3 filtrate; Fgra15-filtrate = *F. graminearum* chemotype 15 filtrate; Fpro-filtrate = *F. proliferatum* filtrate; Fspo-filtrate = *F. sporotrichioides* filtrate; and water = control. Day of incubations were: 1d suspension = spores suspended for 1d in different suspension treatments; 1d sus+PDA = spores suspended for 1d in suspension treatment and then inoculated onto PDA medium for an additional day; 3d suspension = spores suspended for 3d in different suspension treatments; and 3d sus+PDA = spores suspended for 3d in suspension treatment and then inoculated onto PDA medium for an additional day;

Figure 19 are charts showing linear mycelial growths of (A) *Fusarium graminearum* chemotype 3, and (B) *Fusarium graminearum* chemotype 15 in dual-culture assays challenged with *Sphaerodes mycoparasitica* in co-inoculation (same day) (-▲-) and pre-inoculation (1 day prior to *Fusarium* inoculation) (··■··) treatments as well as control (without mycoparasite) (—◆—) for 5 days;

Figure 20 are micrographs of interactions between *F. graminearum* 3-ADON (Fgra3) and 15-ADON (Fgra15) chemotypes on slide culture assays with *S. mycoparasitica* (S) biotrophic mycoparasitic fungus: (A) Single mycelium of *S. mycoparasitica*, (B) *F. graminearum* mycelium with red complex, (C) Absorption of red complex from *F. graminearum* by *S. mycoparasitica* (arrow), (D) Excretion of red complex in crystal-forms (arrows) by *S. mycoparasitica* from mycelium interacting with *F. graminearum* chemotype 3 only, (E) Formation of series of hook-like structures by *S. mycoparasitica*, (F) Parasitism of *F. graminearum* mycelium by *S. mycoparasitica* with formation of hook-shaped structures, excretion of crystal-like compounds (arrows), and internal haustorium, (G) Initiation of penetration-peg formation by *S. mycoparasitica* on *F. graminearum*, (H) Infected or penetrated and non-infected mycelial cells, (I) Branching of haustorium inside *Fusarium* host, (J) Formation of extensive short branching structures by *F. graminearum* chemotype 15 only at the contact zone with *S. mycoparasitica*. Bar scales: (A) to (I) in 5µm, and (J) in 20µm;

Figure 21 is a chart showing the differences in diameters between infected and non-infected *F. graminearum* chemotype 3 (Fgra3) and 15 (Fgra15) host cells in presence of *Sphaerodes mycoparasitica* on slide culture assays. Bars demarked by different lowercase letters represent significant difference in size of infected vs. non-infected hyphae for the two different *F. graminearum* chemotypes at  $P = 0.05$  using a T-test;

Figure 22 are charts of the standard curves of *Fusarium graminearum* chemotype 3 and 15 genomic DNA concentration standards versus cycle threshold (Ct) with PCR reactions performed in triplicate using primer sets; (A) *Tox5-1/2*, with genomic DNA ranging from 270 ng ( $\text{Log}_{10} = 2.90$ ) to 0.27 ng ( $\text{Log}_{10} = -0.60$ ), readings at 0.005 fluorescence line; (B) *Tox5-1/2*, with genomic DNA ranging from 30 ng ( $\text{Log}_{10} = 1.48$ ) to 0.03 ng ( $\text{Log}_{10} = -1.52$ ), readings at 0.005 fluorescence line; and (C) *Fg16NF/R*, with DNA template ranging from 270 ng ( $\text{Log}_{10} = 2.43$ ) to 0.027 ng ( $\text{Log}_{10} = -1.57$ ); in 10-fold dilution series, readings at 0.025 fluorescence line. Error bars indicate standard deviation for the mean of *F. graminearum* chemotype 3 and 15 standard curves derived from *tri5* gene and *F. graminearum* specific primer set;

Figure 23 is a schematic illustration of the experimental set-up for dual-culture assays used to acquire *F. graminearum* chemotype 3-ADON or 15-ADON mycelial plugs for DNAs extraction. The sampling zone (S-zone) indicates the 0.5 x 1.5 cm<sup>2</sup> sample area situated approximately 0.2 cm behind the interaction zone (I-zone). The I-zone represents the interaction or contact zone between *F. graminearum* (Fgra) and *S. mycoparasitica* (SM);

Figure 24 is a chart showing real-time fluorescence curves of *F. graminearum* sequences amplified using *Fg16NF/R* primer set from total DNA extracted from dual-culture assays of *F. graminearum* strains and pre-inoculated *Sphaerodes mycoparasitica* (SM), or singly grown *F. graminearum* cultures of chemotype 3 and 15;

Figure 25 is a chart showing comparisons between different concentrations of DNA from *F. graminearum* chemotype 3 and 15 amplified with *Tox5-1/2* (*tri5* gene specific) and *Fg16NF/R* (*F. graminearum*-specific) primer sets. Fungal DNAs were extracted from 5 d dual-culture assays pre-inoculated with *S. mycoparasitica* for 1 d. With T-test at  $P = 0.05$ , for the comparison between *S. mycoparasitica* treated and non-treated *F. graminearum* chemotype 3 and 15 for

*Tox5-1/2* and *Fg16NF/R* primer sets, respectively. (Log10 transformed for DNA amplified with *Tox5-1/2* primers);

Figure 26 are micrographs of *Fusarium graminearum* mycelia at the contact zone with biological and chemical agents: (A) No visible cell changes with *Sphaerodes mycoparasitica* biotrophic mycoparasite; (B) Cell abruption with *Trichoderma harzianum* necrotrophic mycoparasite; (C) 3-ADON chlamydospores formation in chains when challenged with fungicide; (D) 15-ADON chlamydospores formation in clusters when challenged with fungicide. Scale bars indicate: (A) = 5  $\mu\text{m}$  and (B) – (D) = 10  $\mu\text{m}$ ;

Figure 27 are charts showing gene expression of different *Tri* genes for *F. graminearum* chemotype 3 (3-ADON producer) and *F. graminearum* chemotype 15 (15-ADON producer) in *in vitro* assays with three separate treatments: (A) *Tri4* gene; (B) *Tri5* gene; (C) *Tri6* gene; and (D) *Tri10* gene. Legends: B3 = *S. mycoparasitica* + 3-ADON producing *F. graminearum*; F3 = Folicur + 3-ADON producing *F. graminearum*; T3 = *T. harzianum* + 3-ADON producing *F. graminearum*; B15 = *S. mycoparasitica* + 15-ADON producing *F. graminearum*; F15 = Folicur + 15-ADON producing *F. graminearum*; and T15 = *T. harzianum* + 15-ADON producing *F. graminearum*. Means for three different treatments in *F. graminearum* chemotype 3 and 15 were analyzed separately with LSD test at  $P = 0.05$ . Values are means  $\pm$  SE of three samples;

Figure 28 are charts showing gene expression of *PKS* genes for *F. graminearum* chemotype 3 (3-ADON producer) and *F. graminearum* chemotype 15 (15-ADON producer) in *in vitro* assays with three separate treatments: (A) *PKS4* and (B) *PKS13*. Legends: B3 = *S. mycoparasitica* + 3-ADON producing *F. graminearum*; F3 = Folicur + 3-ADON producing *F. graminearum*; T3 = *T. harzianum* + 3-ADON producing *F. graminearum*; B15 = *S. mycoparasitica* + 15-ADON producing *F. graminearum*; F15 = Folicur + 15-ADON producing *F. graminearum*; and T15 = *T. harzianum* + 15-ADON producing *F. graminearum*. Means for three different treatments in *F. graminearum* chemotype 3 and 15 were analyzed separately with LSD test at  $P = 0.05$ . Values are means  $\pm$  SE of three samples;

Figure 29 is a micrograph of thin liquid chromatography (TLC) analysis for zearalenone (ZEA) extracted from six separate treatments. Legends: B3 = *S. mycoparasitica* + 3-ADON producing *F. graminearum*; F3 = Folicur + 3-ADON producing *F. graminearum*; T3 = *T.*

*harzianum* + 3-ADON producing *F. graminearum*; F15 = Folicur + 15-ADON producing *F. graminearum*; T15 = *T. harzianum* + 15-ADON producing *F. graminearum*, B15 = *S. mycoparasitica* + 15-ADON producing *F. graminearum*. and ZEA = Standard of zearalenone;

Figure 30 is a chart showing the ratio of *F. graminearum* chemotype challenged with Folicur fungicide to *F. graminearum* alone control produced for all four different mycotoxins – ZEA, DON, 3ADON and 15ADON. Legends indicate *F. graminearum* 3-ADON chemotype with Folicur (■) and *F. graminearum* 15-ADON chemotype with Folicur (□);

Figure 31 is a chart showing relative AUR gene expression in *Fusarium* strains after co-culturing with biological and chemical agents. Legend: 15ADON - *F. graminearum*15-*acetyl*-deoxynivalenol chemotype; 3ADON - *F. graminearum*15-*acetyl*-deoxynivalenol chemotype; F.cul. - *F. culmorum*; F.ave. - *F. avenaceum*; B - *Sphaerodes mycoparasitica* (biotrophic mycoparasite); Tricho - *Trichoderma harzianum* (necrotrophic mycoparasite); Fol - Folicur (tebuconazole) fungicide;

Figure 32 is a chart showing changes in AUR gene expression evidenced by the color of fungal hyphae. Legend: Ds #71232B: Red and highly virulent, tolerant at 80<sup>0</sup>C for 4 hours; Es #4E040B: Moderately red and moderately virulent, tolerant 40<sup>0</sup>C for 4 hours; Bs #A86608: White and non virulent, susceptible 40<sup>0</sup>C for 4 hours;

Figure 33 are charts showing the effects of inoculating a *Fusarium*-susceptible barley cultivar with *F. graminearum*, *S. mycoparasitica*, and treating *F. graminearum*-infected barley with different concentrations of *S. mycoparasitica* on: (A) height of the plants; (b) average number of spikes per plant; and (C) the average weight of 5 spikes;

Figure 34 are charts showing the effects of inoculating a *Fusarium*-susceptible wheat cultivar with *F. graminearum*, *S. mycoparasitica*, and treating *F. graminearum*-infected wheat with different concentrations of *S. mycoparasitica* on: (A) height of the plants; (B) average number of spikes per plant; and (C) the average weight of 5 spikes;

Figure 35 is a chart comparing the biocontrol effects of *S. mycoparasitica* on the severity of *Fusarium* head blight symptoms with the protection provided by a commercial fungicide;

Figure 36 are charts showing standard curves of *Fusarium graminearum* chemotype 3 genomic DNA concentration standards versus cycle threshold (Ct) with PCR reactions performed in triplicate using primer sets: (A) *Tox5-I/2*, with genomic DNA ranging from 270 ng ( $\text{Log}_{10} = 2.90$ ) to 0.27 ng ( $\text{Log}_{10} = -0.60$ ), readings at 0.005 fluorescence line; and (B) *Fg16NF/R*, with DNA template ranging from 270 ng ( $\text{Log}_{10} = 2.43$ ) to 0.027 ng ( $\text{Log}_{10} = -1.57$ ); in 10-fold dilution series, readings at 0.025 fluorescence line;

Figure 37 is a chart showing the effects of *S. mycoparasitica* (B) and Folicur fungicide (Fol) treatments on *F. graminearum* chemotype 3-ADON genomic DNA detected in barley spikes employing RT-PCR. Treatments were: Fus – *F. graminearum*; B-Fus – *S. mycoparasitica* with *F. graminearum*; Fol-Fus – Folicur fungicide with *F. graminearum*;

Figure 38 are micrographs showing *Sphaerodes mycoparasitica*–*Fusarium* spp. mycoparasitism assays: (A–a). Hook-shaped contact structures (arrows); (B–b). Clamp-like clasping cells (arrows). “a” and “b” are diagrammatic drawings for (A) and (B) respectively. Scale bars = 5 $\mu\text{m}$ ;

Figure 39 are micrographs showing intracellular parasitism, hyphal inhibition response, and anamorphic stages during the *Sphaerodes mycoparasitica*–*Fusarium* spp. interactions: (A) Intracellular parasitism by *S. mycoparasitica* in *F. equiseti* (arrow); (B) *Fusarium* hyphal inhibition response when challenged with *S. mycoparasitica*; deformation of hyphae into rosette-like shapes (arrow); (C) Hyaline *S. mycoparasitica* anamorphic stages; (D) *Sphaerodes mycoparasitica* anamorphic stages with adsorption of red pigments from *F. culmorum*. Scale bars A, C, D = 5 $\mu\text{m}$ ; B = 20 $\mu\text{m}$ ;

Figures 40A and 40B are micrographs of intracellular parasitism by *Sphaerodes* inside *F. equiseti* (arrows), and 40C and 40D are micrographs of intracellular hyphae produced by *Sphaerodes* inside *F. equiseti* with hook-shaped contact structure (arrows). 40A and 40B were captured under light microscopy; whereas in 40C and 40D hyphae were stained with lactofuchsin and images were captured under fluorescent and confocal laser microscopy, respectively. Scale bars = 5 $\mu\text{m}$ ; and

Figure 41 is a chart showing average hyphal diameters of parasitized and non-parasitized *F. equiseti* cells (■) and *F. culmorum* (□) on 1-week slide-cultures with *Sphaerodes mycoparasitica* biotrophic mycoparasite. Data are means and standard deviations. Same lowercase letters indicate no significant difference between parasitized and non-parasitized hyphae at  $P = 0.05$ , with T-test.

#### DETAILED DESCRIPTION OF THE INVENTION

*Sphaerodes mycoparasitica* (Ascomycetes, Melanosporales), has been isolated from isolates of *Fusarium avenaceum*, *Fusarium graminearum* and *Fusarium oxysporum* originating from wheat or asparagus fields. The species is characterized by a unique combination of ascospore size, shape (fusiform and triangular) and wall ornamentation (reticulate and smooth). Also, conidia are produced from simple phialides on the surface of ascoma peridial wall, on ascoma surrounding hyphae, and on irregularly branched conidiophores arising from hyphae. *S. mycoparasitica* has a phialidic anamorph and produces simple phialides on the surface of ascoma peridial wall or scattered irregularly on ascoma surrounding the hyphae, and on conidiospores. *S. mycoparasitica* forms hook-like structures parasitizing living hyphae of *Fusarium*.

The 1266bp DNA sequence from the large subunit ribosomal RNA gene (LSU) of *S. mycoparasitica* is given in SEQ ID NO: 1.

**SEQ ID NO: 1.**

1 atagggagaa gaagcactgc gattgccta gtaacggcga gtgaagcggc agcagcccag  
61 atttgaatc tggctctttt ggggcccagag ttgtaatctg cagaggaagc gtctggtgcg  
121 gtgccggcct agttccctgg aacgggacgc cgtagagggt gacagccccg tacggtcggc  
181 caccaaacct gtgtgtcgtc ccttcgaaga gtcgcgtagt ttgggaatgc tgcgtaaagt  
241 gggaggatg ctctcctaa ggctaaatac cggccagaga ccatagcgc acaagtagag  
301 tgategaaag atgaaaagca ccttgaatat ggggttaaaa agtacgtgaa attgccaag  
361 gggaagcgtc cgtggccaga ctctgctctt atggatcctc cggctatttc gccggtgcac  
421 tccattagge tggggccagc gtcggtcggc gccggtacta aaagacagcg cgaacgtggc  
481 tctctcggg gagtgttata gcgcgctgtg taatgtgctg gcgccgtccg aggaccgcgc  
541 atttatcaa ggacgtggc gtaatggcca ctacgaccc gtcttgaac acggaccaag  
601 gagtcgcca gagacgcgag tgtgcgggtg acaaaccct gcgcgaagtg aaagcgaacg  
661 ctggtgggaa cctcacggg tgcaccaccg accgatcctg atgtctcgg atggattga  
721 gtatgagcgt ttctggtcgg acccgaaaga ggggtgaacta tgcttgggta ggggtgaagcc  
781 agaggaaact ctggtggagg ccccgtttgg gttctgacgt gcaaatgat ccataaacct  
841 gggcatagcg gcgaaagact aatcgaacct tctagtagct ggttcgcatt ctctctctc  
901 cacgagagag agaaaacctc tgtgatca cgattatcag tgaaaaccac accgagacc  
961 aacggagttc ttctggattt cctcatgctt caattaccac gcctagtgga cctacctgga  
1021 gcgctacaat aaagtcatac gaaaatctcg aagatcgggg tgacggtgag ggatcctaag  
1081 gttctctcgt tgagtgcgtt ggacgggcat ggccgtcagc gatctggggc gaccgttgc  
1141 ggatcataag ggcttttagt cttaggctat tggattgag gggctctgaag acggtaatct  
1201 gaaaccaaag gctttattct aaaccgcgc agcatgggcg tagtaggaag agacagcgaa  
1261 gtctag

The 1233bp DNA sequence from the small subunit ribosomal RNA gene (SSU) of *S. mycoparasitica* is given in SEQ ID NO: 2.

**SEQ ID NO: 2**

```

1      agtgcggcat gttgtagcct aagcaattat acagcgaaac tgcgaatggc tcattatata
61     agttatcgtt tattttagatag tgccttacta cttggataac cgtggtaatt ctagagctaa
121    tacatgctga aagccccgac ttacggaggg gcgtatttat tagattaaaa accaatgccc
181    ttcggggctc tttggtgatt catgataact tctcgaatcg cacggccttg cgccggcgat
241    ggttcattca aatttcttcc ctatcaactt tcgatgtttg ggtagtggcc aaacatgggt
301    gcaacgggta acggaggggt agggctcgac cccggagaag gagcctgaga aacggctcct
361    acatccaagg aaggcagcag gcgcgcaaat tacccaatct caactcgagg aggtagtgac
421    aataaatacc gatgcagggc tctttagggt cttgcaattg gaatgagtac aatttaaact
481    ccttaacgag gaacaattgg agggcaagtc tggtgccagc agccgcggtg actccagctc
541    caatagcgtg tattaaagtt gttgtggta aaaagctcgt agttgaacct tgggcctggc
601    cggtgtgccc gcctaacagc gtgactggt gcggccgggt cttcccaccg cggagccgca
661    tgtcctcac tggcggtgtc ggggaagcgg tacttttact gtgaaaaaat tagagtgctc
721    taagcaggcc tatgctcgaa tacattagca tggaataata gaataggaca gtcgttctat
781    tttgttggtt tctaggacgt ctgtaatgat taacagaaac aatcgggggc gtcagtattg
841    catcgtcaga ggtgaaattc ttgatcgtat gcaagactaa ctactcgaa agcattcgcc
901    aagggtgttt tcattaatca ggaacgaaag ttaggggatc gaagacgatac agataccgctc
961    gtagtcttaa ccataaacta tgccgactag ggatcgggcg gtgtaatttt gaccgctcg
1021   gcacttacga gaaatcttaa gtgcttgggc tccaggggag tatggtcgca aggctgaaac
1081   ttaaagaaat gacgaagggc accaccaagg gtgaacctgc ggcctagttg actcacacgg
1141   gaaactcacg aggtaggaat atgtagatga cggatggagc cttcagaata catatgggca
1201   tgcgctctta ataccggtat tgaaaatggg cag

```

Sample cultures of *Sphaerodes mycoparasitica* have been deposited with Saskatchewan Microbial Collection and Database under the accession number SMCD2220-01, and in the



International Depositary Authority of Canada Collection (1015 Arlington Street, Winnipeg, Canada, R3E 3R2) under the accession number IDAC301008-01.

*Sphaerodes mycoparasitica* may be useful as an anti-fungal agent. *S. mycoparasitica* may be used to treat, ameliorate, or otherwise control infections of *Fusarium* fungi. *S. mycoparasitica* seems particularly useful for treating, ameliorating or otherwise control infections of *Fusarium avenaceum*, *Fusarium graminearum*, and/or *Fusarium oxysporum* and improving plant health or growth.

*S. mycoparasitica* may be used as a prophylactic agent to diminish the chance of a fungal infection, particularly a *Fusarium* infection, from occurring.

*S. mycoparasitica* may be used for treating plants affected by *Fusarium* Wilt Disease or *Fusarium* Head Blight. *S. mycoparasitica* may be used for hindering such conditions from spreading or as a prophylactic measure to diminish the risk of such conditions from occurring.

In the present invention, *S. mycoparasitica* may be isolated from suitable sources. For example, *S. mycoparasitica* may be isolated from *F. graminearum* or *F. avenaceum* isolates originating from wheat fields by the method described in Vujanovic V, *et. al.* Can. J. Microbiol. 48(9): 841-847 (2002). *S. mycoparasitica* is also available deposited with International Depositary Authority of Canada (IDAC301008-01), originally deposited October 30, 2008.

*S. mycoparasitica* may be used to treat, ameliorate, or otherwise control infections of *Fusarium* fungi. *S. mycoparasitica* may be applied in any suitable manner to the organism in need of treatment. For example, *S. mycoparasitica* isolates may be used directly or they may be further processed into soil-applied granular-based, peat-based, seed-applied, or anthesis-applied liquid-based inoculants.

*S. mycoparasitica* may be produced via fermentation and formulated into a pesticide composition. A suitable fermentation process includes, selection of fermentation medium (solid state or submerged), concentration of fermentation constituents, oxygen transfer, incubation temperature, time of harvest and post harvest treatments. The aim is to establish the optimum conditions to ensure an abundant, stable, and efficacious mycoparasitic, microbial population. When formulating *S. mycoparasitica* into a pesticide composition, care should be taken to

selected ingredients that: (i) ensure stability during production, processing and storage, (ii) assist application, (iii) protect the pesticide from unfavourable environmental conditions, and (iv) promote pesticidal activity at the target. Exposure to inactivated host/pathogen or its compounds may improve the activity of *S. mycoparasitica*.

The formulations disclosed herein will usually comprise (i) an active ingredient, (ii) carriers - often an inert material used to support and deliver the densely populated active ingredient to the target, and optionally (iii) adjuvants - compounds that: promote and sustain the function of the active ingredient by protection from UV radiation, that ensure rain fastness on the target, and that retain moisture or protect against desiccation, or promote the spread and dispersal of the biopesticide using standard agriculture equipments such as those disclosed by Hynes and Boyetchko (2006, Soil Biology & Biochemistry 38: 845-84).

Internal Transcribed Spacer ribosomal DNA from *S. mycoparasitica* was sequenced. A 560bp DNA sequence from the ITS region is given in SEQ ID NO: 3.

### SEQ ID NO: 3

```

1      ttcgtttett atgcattgg tgaccagegg agggtcatta cgaatcggac catttatgtc
61     atggctctgc caaccctgtg aactttatac ttgtacgttg cctcggcgga acctgccttt
121    tcggcaggcc gccggccggc atatacgeaa acgctctgaa aaagctccgc gctctatctg
181    aataataaaa cttaacgag taaaaacttt tggcaacgga tctcttggct ctggcatcga
241    tgaaaaacgc agcgaaatgc gatacgtaat gtgaattgca gaattcagtg aaccatcgaa
301    tctttgaacg caccttgccg cggccggtaa tccggcggcc atgcccgtcc gagcgtcgtt
361    tccaccctcg ggagttctcc tectaagaaa atttctcccg gccttgggcc agcgcgttgc
421    gcggtgccc gaccaacggc ggcaggaccg gcgatgtcct ctgtccctg catttatata
481    aaactcgc attggtccccg taaggcttgc cttgcaacca acttctttag gtcgacctca
541    gatcggatag ggatacccgc

```

The present invention provides a primer set, SmyITSF/R (SmyITSF is SEQ ID NO: 4 and SmyITSR is SEQ ID NO: 5), useful for identifying *S. mycoparasitica*.

SEQ ID NO:4 5'-TCATGGCTCTGCCAACCTGTAA-3'

SEQ ID NO: 5 5'-AATGCAGGGCACAGAGGACATCG-3'

The present primer is selective for *S. mycoparasitica* and can be used to assess and quantify the fungus in industrial products, plant materials, seed samples, and environmental samples by using PCR and RT-PCR technologies.

The SmyITSF/R primer set can be used for quantitative real-time PCR technology for analyzing gene expression, in fungal pathogens detection, and in quantification of fungi in living plants.

The SmyITSF/R primer set was tested with SMCD2220-01, seven *Fusarium* species, nine different ascomycetous fungal isolates, two zygomycete fungi, and three basidiomycetous fungal strains.

In PCR, this primer set only amplified SMCD2220-01, not the other fungi. Root biomass, total biomass, root length, total length, and seed germination of *F. graminearum* infected spring wheat were significantly increased with the treatments of *S. mycoparasitica*, as compared to inoculation with *F. graminearum*. In further RT-PCR studies, SmyITSF/R specific primer was used for *S. mycoparasitica* in combination with *F. graminearum*-Fg12NF/R and *Trichoderma*-TGP4-F/R as a control, within which the primer showed high accuracy in assessing biocontrol-pathogen-plant interactions.

Total extracellular protein extracts from *Sphaerodes mycoparasitica* demonstrated significant inhibition of *Fusarium* spore germination. The present invention provides an antifungal agent comprising extracellular protein extracts from *S. mycoparasitica*. Two proteins were isolated from the total protein extract, one having a molecular weight of 13 kDa and one having a molecular weight of 50 kDa. The present invention provides an antifungal agent comprising one or both of these proteins.

Conventional molecular biological techniques may be used to isolate, characterize and produce the present proteins. For example, in order to identify the gene(s) for the present proteins, *S. mycoparasitica* could be challenged with *F. oxysporum* filtrates and upregulated mRNA isolated by standard Northern Blot. cDNA can be produced from the mRNA by Reverse Transcriptase PCR (RT-PCR). The cDNA can be amplified, purified and inserted into an appropriate vector. This vector may be inserted into an appropriate host cell.

Cloning and expression may focus on isolation of genes coding for antimicrobial proteins by designing primers for identified proteins. Isolated genes may be cloned in suitable expression vectors (yeast or bacteria) with suitable/efficient promoters in order to generate industrial strains for large-scale production of the concerned proteins. These vectors can be purchased from Promega, Invitrogen, Clontech, or other companies. The cloned genes may be tested for their protein expression, and the expressed proteins will be purified (using His-tag or other columns). Purified proteins may be tested against plant-pathogenic fungi for their antifungal activity by disc-plate assay and/or microtitre plate assay methods (e.g., Drummond and Waigh, 2000, Recent Research Developments in Phytochemistry.4:143-152). Then, rDNA technologies, involving gene mutation or addition of enhancers, introns or protein-specific promoters (GA inducible), or addition deletion may be used to enhance the production of proteins.

Selected genes may be transformed into wheat and barley to create a transgenic lines with antifungal activity against pathogenic fungi using techniques known to those skilled in these arts (e.g., Sanghyun et.al., 200,8 J. Expt. Botany, 59, 2371-2378; Dennis et. al., 2007. Plant Cell Reports. 26: 631-639).

*S. mycoparasitica* has demonstrated an ability to degrade the deoxynivalenol (DON) mycotoxin. The present invention provides a method of degrading acetyldeoxynivalenol, especially 3-acetyldeoxynivalenol.

*S. mycoparasitica* could be challenged with pure DON and upregulated mRNA isolated by standard Northern Blot. Reverse Transcriptase PCR could be used to identify the gene(s) for the upregulated proteins, an appropriate cDNA construct could then be inserted into an appropriate vector for protein production. Further, recombinant DNA Technology could be applied to create transgenic plants with antimycotoxin properties.

The present invention provides antifungal compositions comprising *S. mycoparasitica*, isolates, cultures, or proteins thereof. The present invention also provides a method of controlling a fungal disease of a plant, and a method for mycotoxin detoxification, which methods comprise applying to the locus of the plant *S. mycoparasitica*, isolates, cultures, or proteins thereof, collectively hereinafter referred to as the 'antifungal agent'.

The compositions of the invention may, for example, be applied to the seeds of the plants, to the growth medium (e.g. soil or water), or to the foliage of the plants.

The present invention provides a composition comprising the antifungal agent and an agriculturally acceptable carrier or diluent, which will ensure stability and performance of the final product. Carrier or diluent should be compatible with the active ingredient, agriculturally acceptable, have a good absorptive capacity and a suitable bulk density, allowing easy particle dispersion and attachment.

The compositions herein may be applied, as in aqueous sprays, granules and dust formulations in accordance with established practice in the art. An aqueous spray is usually prepared by mixing a wettable powder or emulsifiable concentrate formulation of a compound of the invention with a relatively large amount of water to form a dispersion.

Wettable powders may comprise a finely divided mixture of the antifungal agent, a solid carrier, and a surface-active agent. The solid carrier is usually chosen from among attapulgite clays, kaolin clays, montmorillonite clays, diatomaceous earths, finely divided silica, purified silicates, or combinations thereof. Surfactants which may be useful herein have wetting, penetrating, and/or dispersing ability. They are typically present in an amount of from about 0.5% to about 10% by weight. Surfactants herein may be chosen from, for example, alkylbenzenesulfonates, alkyl sulfates, naphthalenesulfonates and condensed naphthalenesulfonates, sulfonated lignins, and non-ionic surfactants.

Emulsifiable concentrates may comprise the antifungal agent of the invention in a liquid carrier, the carrier being a mixture of a water-immiscible solvent and a surfactant. Solvents that may be useful herein include aromatic hydrocarbon solvents such as the xylenes,

alkylnaphthalenes, petroleum distillates, terpene solvents, ether-alcohols, organic ester solvents or suitable combinations thereof.

When a composition of the invention is to be applied to plant debris or litter, in order to control of the source of contamination and inoculant dispersion, or to the soil, as for pre-emergence protection, granular formulations or dusts are sometimes more convenient than sprays.

In one embodiment the antifungal agents herein are encapsulated into alginate pellets. The pellets may be prepared in any suitable manner. For example, one useful method is described in Harveson et. al., (2002, Plant Disease; Vol. 86, No. 9 1025-1030).

The composition may comprise other active substances useful in an antifungal agent. For example, the compositions herein may comprise other antifungal agents such as *Trichoderma*, sulfur, neem oil, rosemary oil, jojoba oil, *Bacillus subtilis*, allylamines (e.g. terbinafine, antimetabolites (e.g. flucytosine), azoles (e.g. ketoconazole, itraconazole), echinocandins (e.g. caspofungin), polyenes (e.g. amphotericin B), systemic agents (e.g. griseofluvin), or combinations thereof.

The compositions herein may contain from about 0.1% to about 95%, by weight, of the antifungal agent and from about 0.1% to about 95%, by weight, of the carrier and/or surfactant. The direct application to plant seeds prior to planting may be accomplished in some instances by mixing either a powdered solid compound of the invention or a dust formulation with seed to obtain a substantially uniform coating which is very thin and represents only one or two percent by weight or less, based on the weight of the seed. In some instances, however, a non-phytotoxic solvent such as methanol is conveniently employed as a carrier to facilitate the uniform distribution of the compound of the invention on the surface of the seed.

The compositions herein may be useful in the treatment of fungal infections in plants. Consequently, the compositions herein may comprise *S. mycoparasitica*, isolates, cultures, or proteins thereof and a pharmaceutically acceptable carrier.

*S. mycoparasitica* sporulates in the presence of *Fusarium*. *S. mycoparasitica* ascospores proved resistant to germination under different standard laboratory conditions (sterile distilled

water, on water agar and commercially available media) and heat or cold-shock treatments. In contrast, spore germination was obtained on general potato dextrose agar medium amended with *Fusarium*-filtrates. Significant improvement in percentage of spore germinations were obtained for the spores suspended in *Fusarium*-filtrates. *F. avenaceum* and *F. oxysporum* filtrates induced the highest germination, whereas *F. sporotrichioides* and *F. proliferatum* triggered lower germination frequency. Filtrates of beneficial fungal inoculants: *Trichoderma harzianum* (RootShield<sup>®</sup> available from BioWorks Inc., Victor, NY, USA; RootShield is a registered trademark of BioWorks Inc.), *Penicillium bilaii* (JumpStart<sup>®</sup> available from Novozymes Biologicals Ltd., Saskatoon, SK, CA; JumpStart is a registered trademark of Philom Bios Inc.), and *Chaetomium globosum* had no impact on germination. Ascospores suspended in *F. avenaceum*-filtrate, showed double-polar germination pattern. On the other hand, when suspended in *F. oxysporum*-filtrate, significant amount of spores demonstrated a single-polar germination pattern. *S. mycoparasitica* grown on *F. oxysporum* kept the same mycoparasitic germination patterns in three offspring generations when transferred on *F. avenaceum* which indicate a stable genome-regulated expression.

The present invention provides a method of sporulating *S. mycoparasitica* by exposing them to *F. avenaceum* or *F. oxysporum*, or filtrates, extracts, or compositions thereof.

The present invention provides a method for producing an antifungal composition comprising *S. mycoparasitica* or isolates, cultures, genes or proteins thereof. The method comprises:

- (a) inducing *S. mycoparasitica* sporulation;
- (b) culturing *S. mycoparasitica*;
- (c) harvesting *S. mycoparasitica*.

The method may further comprise any of the following optional steps:

- (d) storing/preserving *S. mycoparasitica*;
- (e) applying *S. mycoparasitica*;
- (f) assessing *S. mycoparasitica*;
- (g) producing *S. mycoparasitica* proteins;
- (h) harvesting *S. mycoparasitica* proteins;

- (i) fractionating *S. mycoparasitica* proteins;
- (j) separating *S. mycoparasitica* proteins;
- (k) storing/preserving *S. mycoparasitica* proteins;
- (l) applying *S. mycoparasitica* proteins;
- (m) assessing *S. mycoparasitica* proteins.

## EXAMPLES

### Example 1

#### *Sampling, fungal growth and microscopy*

Myclobutanil-agar (MBA) medium was used for selective isolation of various *Fusarium* taxa and associated biotrophic mycoparasites from Canadian agriculture fields using the method described in Vujanovic V, *et. al.* (2002, Can. J. Microbiol. 48(9): 841-847). *Sphaerodes* was recovered occasionally from *F. graminearum* and abundantly from *F. avenaceum* isolates originating from wheat fields in Saskatchewan; it was also isolated from *Fusarium oxysporum* from asparagus fields in Quebec, Canada. A monosporal, single culture of the mycoparasite was obtained from each *Fusarium* species according to the method proposed by Harveson & Kimbrough (2001, Int. J Plant Sci. 162(2):403-410). Single ascospore isolates were maintained on Potato dextrose agar (PDA) (Difco, BD Biosciences, Mississauga, ON, CA) supplemented with antibiotics (100  $\mu\text{g l}^{-1}$  streptomycin sulphate and 13  $\mu\text{g l}^{-1}$  neomycin sulphate; Sigma-Aldrich Canada Ltd., Oakville, ON, CA) and stored at  $-80^{\circ}\text{C}$  in Saskatchewan Microbial Collection and Database (SMCD2220-01) and in the International Depository Authority of Canada (IDAC301008-01) collections. Fungal growth was assessed on modified Leonian's agar (MLA) and Potato dextrose agar (PDA) media. Biotrophic interactions between *Sphaerodes* and *Fusarium* strains were examined with the slide culture method proposed by Jordan & Barnett (1978, *Mycologia* 70(2):300-312). Morphological studies of ascomata, ascospores, mycelia, and anamorphic structures were performed after two weeks of incubation ( $21^{\circ}\text{C}$  -  $22^{\circ}\text{C}$ ) under a Carl Zeiss Axioskop2™ with a Carl Zeiss AxioCam ICcl™ camera. Fungal materials for microscopic observation were mounted in lactofuchsin and lactophenol cotton blue dyes.



### *DNA extraction, amplification and sequencing*

Three *Sphaerodes* strains: SMCD 2220-01 on *F. avenaceum* from wheat, SMCD 2220-02 on *F. graminearum* from wheat, and SMCD 2220-03 on *F. oxysporum* from asparagus were cultured on PDA medium at 21°C for a week prior to DNA extraction. Genomic DNA was extracted with the DNeasy<sup>®</sup> Plant Mini Kit (Qiagen Inc., Mississauga, ON, CA; DNeasy is a registered trademark of Qiagen GmbH Corp, Hilden, Fed. Rep. Germany). LSU (large subunit) rDNA fragments were amplified using primer sets NS1/NS6 using techniques known to those skilled in these arts (e.g., Gardes & Bruns, 1993, *Molecular Ecology* 2: 113-118; White *et al.* 1990, *PCR Protocols: a guide to methods and application*: 315-322. Academic Press, New York) and LS1/LR5 (e.g., Hausner *et al.* 1993, *Canadian Journal of Botany* 71: 52-63; Rehner & Samuels, 1995, *Canadian Journal of Botany* 73 (Suppl. 1): S816-S823; Zhang & Blackwell, 2002, *Mycological Research* 106: 148-155). Target regions of fungal genomic DNA samples were amplified using polymerase chain reaction (PCR) in a 25µl reaction mixture containing 2.5µl of 10X buffer, 5µl of Q buffer, 0.5µl 10mM dNTPs, 1µl of each primer, 0.13µl of 0.625 unit of Taq DNA Polymerase, 2µl of extracted fungal DNA, and 12.87µl of sterilized ultra-pure Millipore<sup>™</sup> water. The Qiagen Taq PCR core kits<sup>™</sup> were purchased from Qiagen Inc., Mississauga, ON, CA. Purified DNA PCR products were sequenced.

### *Sequence alignment and phylogenetic analyses*

Sequences of LSU from this study and sequences retrieved from GenBank were aligned using Clustal X software (version 1.82) (Thompson *et al.* 1997, *Nucleic Acids Research* 24: 4876-4882), and edited in Bioedit (Hall, 1999, *Nucleic Acids Symposium Series* 41: 95-98). Distance trees were produced with PAUP (Phylogenetic Analysis Using Parsimony) 4.0b10 software (Swofford 2000) using a neighbor-joining approach, and validated using bootstrap analyses with 1,000 repetitions. A fungal distance tree was prepared with sequences showing bootstrap values higher than 50%. Trees were rooted with sequences *Xylaria hypoxylon* U47841.

Taxonomy: *Sphaerodes mycoparasitica* Vujanovic, sp. nov. (Figures 1-5) [MycoBank no: MB 515144], in the International Depository Authority of Canada as *Sphaerodes mycoparasitica* strain IDAC 301008-01.

*Coloniae* in agaro potato dextrosum lentior crescents, 4.0 cm ad 7d, floccose, pallido-brunneis. *Hyphis* septatis, ramosis, anastomosantibus, laevibus, palide fulvis, 2.5-5.0  $\mu\text{m}$  diam compositum. *Ascomata* superficialia vel immersa, pyriformia vel globosa, ostiolata, flavo-brunnea, 250-300  $\mu\text{m}$  longa, 200-280 $\mu\text{m}$  diam. *Collum* nul, conicum vel cylindricum, 30-75 $\mu\text{m}$  longum, (0-) 50-70  $\mu\text{m}$  latum ad basim. *Peridium* membranaceum, cellulis 8-15  $\mu\text{m}$ , e 3-6 stratis, 8-15  $\mu\text{m}$  crassum, textura angulari compositum. *Setae* rectae vel parum curvae, hyalinae vel dilute flavae, crassitunicatae, 10-40  $\mu\text{m}$  longae, septatae. *Asci* 8-sporei, ovoidei vel clavati, 50-75 X 17-25  $\mu\text{m}$ , superne late rotundati, brevistipitati, tenuitunicati, evanescentes. *Paraphysis* nullis. *Ascosporeae* unicellulares, irregulariter biseriatae, primum hyalinae, deinde brunneae vel atrobunneae, crassitunicatae, fusiformes, 18-24 X 9-12  $\mu\text{m}$ , reticulatae, costis protrudentibus, e polo visae polygonales, utrinque umbonatae, foramine germinali praeditae. *Phialidis* hyalinis, status conidialis.

*Culture characteristics:* Colonies of *Sphaerodes mycoparasitica* strain IDAC 301008-01 cultured on MLA grew more rapidly than on PDA, 1.1 cm versus 0.6 cm per day (21-22°C), consisting of slightly submerged mycelium and aerial hyphae, granulose due to production of ample number of ascomata. On MLA (Figures 1A and 1C) and PDA Figures 1B and 1D), the cultures produced a woolly mycelium, yellowish to pinky-brownish on both sides. At 37°C, no growth was registered. *Hyphae* were white to pale yellow, 2.5-5.0  $\mu\text{m}$  diam., septate, anastomosis occurred soon after ascospore germination (Figure 2). Colonies on Potato dextrose agar (PDA) spread with abundant, white to pale yellow aerial mycelium and low number of ascomata. Ascomata, perithecial or cleistotecial, scattered or aggregated in small groups, superficial, pyriform to globose, ostiolate (when mature), light to dark yellowish brown, translucent, appearing black due to mass of mature ascospores, 250-300  $\mu\text{m}$  high, 200-280  $\mu\text{m}$  diam. *Neck* absent to short conical or cylindrical 25-75  $\mu\text{m}$  long, 20-70  $\mu\text{m}$  wide at the base, sometimes surrounded with a crown of short, upright setae, 10-40  $\mu\text{m}$  long. *Peridium* membranaceous, 3-6-layered, 8-10  $\mu\text{m}$  thick, translucent, pale yellow to light brown, composed by cells of 8-15  $\mu\text{m}$  diam. disposed in *textura angularis*.. *Asci* 8-spored, clavate, 50-75 X 17-25  $\mu\text{m}$ , rounded at apex, without apical structures, thin-walled and evanescent when mature. *Paraphyses* absent. *Ascospores* irregularly arranged inside the asci, at first hyaline but becoming brown to dark brown, thick-walled, single-celled, fusiform to rarely triangular, 18-24 X 9-12  $\mu\text{m}$ ,

reticulate to rarely smooth, with irregular transverse sections, and with a strongly umbonate germ pore at each end. *Phialides* hyaline, ampulliform produced directly on ascomata or on hyphae surrounding ascomata, and on irregularly branched conidiophores.

*S. mycoparasitica* strain IDAC 301008-01 has a unique combination of features shown in Figure 3. The ascomata height is generally less than 250  $\mu\text{m}$  with a conical to cylindrical neck (Figure 3A). The setae length is generally less than 40  $\mu\text{m}$  (Figure 3B) and the spore length is generally less than 23  $\mu\text{m}$  (Figure 3C). The spores show a conspicuous wall ornamentation and prominent irregular longitudinal ribs indicated by the arrows in Figure 4A. The spores of *Sphaerodes quadrangularis* are shown in Figure 4B for comparison. *S. mycoparasitica* strain IDAC 301008-01 spores occasionally show a triangular shape (Figures 3D and 3E). The formation of starting and mature ascoma of *S. mycoparasitica* strain IDAC 301008-01 are shown in Figure 3H. This strain produces simple phialides on the surface of ascoma peridial walls or alternatively, the phialides may be scattered irregularly on ascoma surrounding the hyphae, and on conidiospores with a distinctive branching pattern (Figures 3F and 3G). *S. mycoparasitica* strain IDAC 301008-01 forms hook-like structures for parasitizing living hyphae of *Fusarium* (Figures 3I and 3J).

### Example 2

*Sphaerodes mycoparasitica* (SMCD 2220-01) 21°C isolates were cultured in potato dextrose broth (PDB) culture media. About 3 ml of culture were transferred to 250 ml Erlenmeyer flasks containing 50 ml PDB growth medium. The flasks were incubated for 7 days on a rotary shaker (150 rpm) at room temperature.

*Extracellular protein extraction:* Young mycelia were filtered through Whatman® No.1 filter paper (Whatman is a registered trademark of Whatman International Ltd., Kent, UK). Filtered culture medium, containing extracellular proteins, and were concentrated by Amicon ultrafiltration centrifuge tube with a 3000 Dalton cut-off membrane by centrifugation at 4000 rpm at 4°C.

*Disc diffusion assay:* Antifungal activities of extracellular protein extracts were tested under sterile conditions by radial disc plate diffusion assay as described by Roberts & Selitrennikoff

(1986, *Biochim. Biophys. Acta*, 880: 161-170). The assay of the isolated protein for antifungal activity toward *F. oxysporum* and *F. graminearum* was carried out in petri plates containing potato dextrose agar. Mycelial plugs from actively growing fungal plates were placed in the center of the petri plates and sterile filter paper discs (5-mm diameter of Whatman<sup>®</sup> filter paper no. 1) were placed on the agar surface at a distance of 0.5 cm away from the rim of the mycelial colony. An aliquot (60  $\mu$ L) containing 2.5  $\mu$ g of extracellular protein was added to a disk. Sterile distilled water and buffer served as controls. The plates were then incubated at room temperature for 4 days and examined for inhibition. The area of the mycelial colony was measured and the inhibition of fungal growth was determined by calculating the % reduction in area of mycelial colony with the controls (Figure 5A). After 4<sup>th</sup> day of germination, about 30% inhibition of the hyphal extension of *F. oxysporum* and 35% inhibition of hyphal extension of *F. graminearum* were observed (Figure 5B)

*Fast protein liquid chromatography (FPLC) of extracellular proteins:* Proteins were fractionated through Superdex 75 GL 10/30 column using FPLC AKTA<sup>®</sup> purifier system (GE Healthcare, Biosciences AB, CA: AKTA is a registered trademark of GE Healthcare Bio-Sciences AB Ltd., Uppsala, Sweden) according to the manufacturer's instructions. The column was previously equilibrated with sterile water and with 50 mM sodium phosphate buffer, pH 7.0 containing 0.15 M NaCl, followed by protein injection (about 500  $\mu$ L) and elution of proteins with the same buffer with flow rate of 1.0 ml/min. Fractions of 0.8 ml were collected in each tube. Upon gel filtration on Superdex<sup>®</sup> 75 (Superdex is a registered trademark of GE Healthcare Bio-Sciences AB Ltd., Uppsala, Sweden), proteins were resolved into two distinct peaks (f1 and f2) and few smaller peaks (Figure 6). Samples from all the peak fractions were pooled, precipitated and tested for their antifungal activity.

*Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE):* SDS-PAGE (12%) of proteins recovered from FPLC fractions was performed according to the method of Laertunli (1970, *Nature* 227: 680-685). All peaks giving FPLC fractions were pooled and recovered by precipitation in 1:4 volume of chilled acetone and kept at -20°C overnight. After centrifugation at 12,000 g for 10 min, precipitated proteins (pellets) were dissolved in minimum amount (30  $\mu$ L) of assay buffer. Proteins were analyzed by SDS-PAGE having 5% stacking gel (pH 6.8) and 12% separating acrylamide gel (pH 8.8) in Tris-glycine buffer (pH 8.3) and appropriate markers.

Prior to SDS-electrophoresis, the protein was mixed with an equal volume of sample buffer (60 mM Tris-HCl buffer, 4% SDS, pH 6.8) containing 5%  $\beta$ -mercaptoethanol. A mixture of standard marker proteins (Bio-Rad protein markers, Bio-Rad Laboratories Inc., Mississauga, ON, CA) was used. V All samples were heated for 5 min at 95°C and cooled to room temperature before loading on gel. Proteins were visualized by silver staining method (Bio-Rad Silver staining kit, Bio-Rad Laboratories Inc., Mississauga, ON, CA). Proteins molecular masses were estimated by comparison with the mobilities of standard molecular mass markers. Protein bands of molecular weight 13 kDa and 50 kDa were detected in f1 and f2 peaks respectively (Figure 7).

*Microtitre plate assay:* Percentage inhibition of spore germination was performed by microtitre plate assay method (Ghosh, 2006, *Ann. Bot.* 98: 1145-1153; Yadav *et al.* 2007, *J. Med. Microbiol.* 56: 637-644) to test the antifungal activity of proteins recovered from FPLC fractions. The possible toxicity of the fractionated proteins was tested by a percentage growth inhibition assay using the *F. oxysporum* and *F. graminearum*. The *in vitro* antifungal activities of fractionated proteins were determined in 96-well microtiter plates. In microplate wells, 10  $\mu$ l of potato dextrose broth (PDB; BD Biosciences, Mississauga, ON, CA) was mixed with 3  $\mu$ l of spore suspensions of *F. oxysporum* and *F. graminearum*. An aliquot of 7  $\mu$ L of different peak containing proteins fractions were added to suspensions in microtitre plates (12-8 wells). Water 25 and buffer were used as negative controls. The microtitre plate was then incubated at room temperature in dark. Observations were made for inhibition in spore germination in both untreated and treated wells after 24 h using inverted and fluorescent microscope. The number of germinated and non-germinated spores and percentage of area covered by mycelia in microscope were used to determine the percentage of growth inhibition. The f1 and f2 protein-containing peaks had inhibitory effects on spore germination after 24 hrs compared to the control treatment.

The discharge of spores from sporangia was inhibited by these protein fractions. Secondary branching (mycelia formation) was observed in control treated spores of *F. oxysporum* (Figure 8A-c) and *F. graminearum* (Figure 8B-c). No branching was observed in f1 and f2 protein-treated spores of *F. oxysporum* (Figures 8A-a-b) and *F. graminearum* (Figure 8B-a-b). The size of germ tubes (hyphae) from control-treated spores was more than 100  $\mu$ m compared to 10-15  $\mu$ m for protein-treated spores (Figures 8A-a-b; 8B-a-b).

### Example 3

Active growing mycelia of *S. mycoparasitica* (SMCD 2220-01) were cultivated on Potato 10 Dextrose Broth (PDB) in a shaking flask at 21°C for 3 days and then washed. Wet mycelium (0.1 g) were resuspended in 1 ml of PDA medium supplemented 100 µg/mL 3-ADON (Sigma-Aldrich Canada Ltd., Oakville, ON, CA) and incubated for 10 days.

A 0.2cm<sup>2</sup> plug *S. mycoparasitica* was incubated in 1ml PDB containing 3-ADON at a 15 concentration of 100 µg at RT for 7 days. The sample was analysed for DON degradation by TLC and HPLC assays.

#### *Extraction of 3-ADON:*

3-ADON was extracted by the method disclosed by Vasavada and Hsieh (1987, Appl. Micro. Biotechnol. 26: 517-521). The spent medium in sampled flasks was filtered through the Whatman<sup>®</sup> filter paper avoiding the mycelia. 3-ADON was extracted from the medium by three 10 ml volumes of ethyl acetate. The mixture of the medium and solvent was vigorously shaken and allowed to stand for 5 min for separation of phases. The organic phase was siphoned off and passed through sodium sulfate to remove residual water. The solvent was allowed to evaporate at room temperature. The residue was redissolved in acetonitrile for analysis.

#### *Analysis of DON by TLC:*

Thin layer chromatography (TLC) was used to analyze the DON. TLC was performed by the method disclosed by Andrea *et.al.* (2004, J. Basic Microbiol. 44: 147-156). Dried residues were dissolved in 75 µl and loaded onto silica gel TLC plates coated with fluorescent indicator (60 F<sub>254</sub>, 0.2 mm layer; Merck Frosst Canada Ltd, Kirkland, PQ, CA). The spots were focused in the solvent system with ethyl acetate-toluene (3:1) for 40 5 min. The gel plates were sprayed with 20% aluminum chloride in 95% ethanol. Trichothecenes were visualized as dark spots under short wavelength UV light (254 nm). Chromatograms were photographed using gel documentation system. A known standard control of pure 3-ADON (Sigma-Aldrich Canada Ltd., Oakville, ON, CA) was employed to compare with other treatments. The sample containing *S. mycoparasitica* showed significantly less 3-ADON than the control sample.

### *High Performance liquid chromatography (HPLC)*

A Waters HPLC with : 250 x 4.60 mm id Prodigy 5 $\mu$  ODS (3) 100A, 5 $\mu$  micron C<sub>18</sub> column (Phenomenex Inc., Torrance, CA USA) and a photodiode-array (PDA) detector was used with a gradient solvent system (water-acetonitrile containing 0.005% (v/v) trifluoroacetic acid). The PDA detector measured the UV spectrum (200-600 nm). Samples were dissolved in acetonitrile and 20  $\mu$ l was loaded onto the column using an automatic injector. The DON was eluted with solvent as a mobile phase at a rate of 1 ml min<sup>-1</sup>. 3-ADON obtained from Sigma-Aldrich (Sigma-Aldrich Canada Ltd., Oakville, ON, CA) was used as a standard. The results are shown in Figure 9. The sample with *S. mycoparasitica* contained a much reduced level of 3-ADON.

### Example 4

Potato dextrose agar (PDA), Potato dextrose broth (PDB), Yeast extract, Malt extract agar (MEA), Agar, and Peptone were purchased from BD Biosciences (Mississauga, ON, CA) . Streptomycin sulphate, Neomycin sulphate, and other reagents of analytic grade were from Sigma-Aldrich Canada Ltd. (Oakville, ON, CA) .

### *Fungal strains and growth conditions*

Four *phytopathogenic Fusarium* strains (*F. avenaceum*, *F. sporotrichioides*, *F. oxysporum*, and *F. proliferatum*); three beneficial fungal inoculants (*Trichoderma harzianum* (RootShield<sup>®</sup>), *Penicillium bilaii* (JumpStart<sup>®</sup>), and *Chaetomium globosum*); and one mycoparasitic fungal strain (*Sphaerodes mycoparasitica* SMCD 2220-01) isolated from *F. oxysporum* host, were maintained on PDA amended with antibiotics (100 mg/L streptomycin sulphate and 12 mg/L neomycin sulphate) and used throughout this study. *S. mycoparasitica* mycoparasitic fungal isolate was separated from their *F. oxysporum* host, and monosporium cultures were achieved according to methods described by Harveson RM, Kimbrough JW, (2001, Int. J. Plant Sci. 162: 403-410) with few modifications. Mature perithecia were picked up, suspended, and shaken in 2-ml sterilized water blanks. The spores-suspension was then spread on PDA plates. After few minutes, with the assistance of a Carl Zeiss Stemi 1000<sup>™</sup> dissecting microscope, individual spores were removed and transferred to PDA supplemented with 100

m1/1 of *Fusarium* filtrate. All fungal isolates were maintained in the culture collection of the Saskatchewan Microbial Collection and Database, Canada (SMCD).

### *Spores production*

Mycelium plugs from the margin of the active growing monosporial *Sphaerodes* culture were cut and placed on Modified Leonian's Agar (MLA: Maltose, 6.25g; Malt extract, 6.25g; KH<sub>2</sub>PO<sub>4</sub>, 1.25g; Yeast extract, 1.0g; MgSO<sub>4</sub>.7H<sub>2</sub>O, 0.625g; Peptone, 0.625g; Agar, 20g, and 1L of dH<sub>2</sub>O. Inoculated MLA plates were incubated at room temperature (23°C) in dark condition for a month before collecting the spores for germination assays. With the assistance of dissecting microscope, mature spores exuded from the ascomata and located on the ostiolar opening were harvested by picking up carefully with sterile needles and added to 2 ml tubes in 1 ml sterilized distilled water. The suspension was then filtered through four thin layers of cheesecloth to remove the remaining vegetative cells or mycelia. The density of the spore suspension was counted with a haemocytometer and adjusted to approximately 5-6x10<sup>5</sup> spores per ml in sterilized water. Freshly prepared aqueous spore suspensions were used.

### *Preparation of the fungal filtrates*

Four pathogenic *Fusarium* strains and three beneficial fungal strains were grown in shake cultures for 7 days at room temperature (23°C) in 500 ml flasks, each with 100 ml PDB medium.

After incubation of 7 days in PDB medium, mycelia were removed by filtering through filter paper and the filtrates were then filter-sterilized. Freshly preparations fungal filtrates were employed for the spore germination tests.

*Sphaerodes mycoparasitica* strain SMCD 2220-01 was found to sporulate on or when inoculated together with *Fusarium* species, such as *F. avenaceum* and *F. oxysporum*. *S. mycoparasitica* was observed to produce approximately the same amount of ascomata on both *F. oxysporum* and *F. avenaceum*. *S. mycoparasitica* was not found to produce fruiting bodies on other *Fusarium* or fungal strains such as *F. proliferatum*, *F. sporotrichioides*, *P. bilaai*, *T. harzianum*, and *C. globosum*. In the contact zone between these biotrophic mycoparasites and *Fusarium* species, hook-shaped contact structures were formed.



### *Effects of heat and cold treatments on Sphaerodes spore germination*

Aqueous ascospore suspensions were heat-shocked at 60°C and 65°C for 20 min, and cold-treated at 4°C, -20°C, and -70°C for 5 min and 20 min. The heat- and cold-shocked spores were then transferred and inoculated onto WA and PDA for 1 day and 3 days. The readings of spore germination were checked daily. Spores not subjected to heat and cold treatments were used as control.

There was no germination observed in any of the heat and cold treatment groups for 3 days. Additional observations were continued up to 7 days and no spore germination was observed.

### *Germination on various media*

Aqueous spore suspensions were transferred and inoculated onto the surface of the following media:

- 1.5% water agar (WA), PDA, MLA, MEA, Carnation leaves agar (CLA);
- 1.5% water agar plus 100 ml/l of *Fusarium* filtrate;
- PDA with 100 ml/l of *Fusarium* filtrate.

*Fusarium* strains utilized were *F. avenaceum*, *F. oxysporum*, *F. proliferatum*, and *F. sporotrichioides*. Ascospore-inoculated plates were then incubated at room temperature (23°C) for 3 days. Spore germination was examined daily.

The results are summarized in Table 1. Each incubation day for *Sphaerodes* was analyzed separately. Numbers in each column represent the mean of ascospore germination (in %) ± standard deviation. Means within each column for each medium treatment followed by the same letter are not significantly different at  $P \leq 0.05$  after Mann-Whitney U test.

Table 1:: Sporulation of *S. mycoparasitica* on various media

Medium	<i>S. mycoparasitica</i>	
	Day 1	Day 3
Water agar (WA)	0 d	0 d
<i>F. avenaceum</i> -WA	11.1 ± 1.7 b	35.2 ± 2.6 b
<i>F. oxysporum</i> -WA	1.6 ± 1.3 c	18 ± 1.4 c
<i>F. proliferatum</i> -W A	0 d	0 d
<i>F. sporotrichioides</i> -WA	0 d	0 d
Potato dextrose agar (PDA)	0 d	0 d
<i>F. avenaceum</i> -PDA	21.5 ± 1.4 a	63.2 ± 2.4 a
<i>F. oxysporum</i> -PDA	2.5 ± 2 c	37.5 ± 2 b
<i>F. proliferatum</i> -PDA	0 d	0 d
<i>F. sporotrichioides</i> -PDA	0 d	0 d

#### *Effects of fungal filtrates on spore germination*

The aqueous spore suspensions were suspended in the filtrates of four separate pathogenic *Fusarium* and three beneficial fungal strains for 1 day and 3 day in the ratio of 1:2 (1 part of aqueous spore suspension: 2 parts of fungal filtrate), and the filtrate-suspended spores were then transferred and inoculated onto PDA for additional 1 day. Control treatments were suspended with sterilized distilled water or PDB.

#### *Spore germination*

Microscopic assessments of ascospore germination were conducted after incubation for 1 day and 3 days. Percentage of germinated spores was obtained by scoring the spores on the Petri dish through utilizing the 200x and 400x objectives of the Carl Zeiss Axioskop2 microscope and systematically choosing 50 spores, starting at the top right corner and continuing to count till 50. Each drop of ascospore suspensions on a medium plate was considered as a subunit, and there were three subunits per plate. Each medium plate for each treatment was replicated, there were three replicas per treatments. The experiments were repeated twice. An ascospore was only considered as germinated spore when the germ tube was visibly noticeable. Germinated ascospores were counted and recorded as a percentage of the total ascospore number.

The results are summarized in Table 2. Numbers in each column represented mean of ascospore germination (in %)  $\pm$  standard deviation. Each incubation day for *Sphaerodes* was analyzed separately. Means within each column of *Sphaerodes* for each filtrate-suspension treatment followed by the same letter are not significantly different at  $P \leq 0.05$  after Mann-Whitney U test.

Table 2: Effects of fungal filtrates on *S. mycoparasitica* spore germination

<i>Sphaerodes</i> sp.	Spore germination (in %)			
	1d suspension	1d sus + 1d PDA	3d suspension	3d sus + 1d PDA
Water	0 c	0 e	0 e	2 $\pm$ 1.2 c
PDB	0 c	0 e	0 e	2.8 $\pm$ 1.9 c
<i>F. avenaceum</i> -filtrate	89.2 $\pm$ 6.2 a	91 $\pm$ 8.4 a	93.2 $\pm$ 6.7 a	94.2 $\pm$ 7.9 a
<i>F. oxysporum</i> -filtrate	0 c	81.8 $\pm$ 11 b	85 $\pm$ 7.9 b	91 $\pm$ 10 a
<i>F. proliferatum</i> -filtrate	2.6 $\pm$ 3.6 b	3.2 $\pm$ 4.6 d	10 $\pm$ 7.4 c	11.2 $\pm$ 8 b
<i>F. sporotrichioides</i> -filtrate	0 c	3 $\pm$ 2.8 d	3.4 $\pm$ 3.3 d	10 $\pm$ 5.6 b
<i>P. bilaii</i> -filtrate	0 c	0 e	0 e	0 c
<i>T. harzianum</i> -filtrate	0 c	9 $\pm$ 1.1 c	1.6 $\pm$ 1.6 d	10.4 $\pm$ 1.5 b
<i>C. globosum</i> -filtrate	0 c	0 e	0 e	0 c

### Example 5

Since *F. graminearum* 3-ADON is the most pathogenic and mycotoxigenic in wheat, 3-ADON was used to quantifying mycoparasite-*Fusarium*-wheat root interactions under Phytotron controlled conditions.

#### *Fungal strains and growth*

*Fusarium* strains: *F. graminearum* 3-ADON strain SMCD2243, biotrophic mycoparasite *Sphaerodes mycoparasitica* SMCD2220-01, and *Trichoderma harzianum* T-22 (RootShield-commercial product) as a control strain with mycoparasitic properties were maintained on PDA amended with antibiotics.

### *Taxon specific primers*

Specific primer sets for: *F. graminearum*-Fgl 2NFIR (Nicholson et al. 1998. *Plant Pathology* 53: 17-37.); *Trichoderma*-TGP4-F IR (Kim and Knudsen 2008. *Applied Soil Ecology* 40: 100-108); and *S. mycoparasitica* SMCD2220-01-SmyITSF/R were used in this study.

### *Mycoparasite/Fusarium/plant interactions*

Wheat CDC-TEAL 2001 plants were inoculated with mycotoxigenic and pathogenic *Fusarium* as well as *Trichoderma* (control mycoparasite) or *Sphaerodes* SMCD2220-01, and were subjected to RT-PCR quantification to assess the amount of fungal DNAs in roots of wheat under controlled conditions after 14 days of incubation in Phytotron conditions.

### *Growth conditions and fungal inoculation*

Five mycelial plugs from *S. mycoparasitica*, *T. harzianum*, and *F. graminearum* were cut, transferred and grown in three separate shake cultures for 14 d at room temperature (23°C) in 500 mL flasks each with 100 mL of PDB (potato dextrose broth). After incubation of 14 d in PDB medium, the fungal cultures were filtered through a Whatman® No. 1 filter paper to remove the liquid medium. The mycelia were then transferred to 50 mL sterile Falcon tubes with 20 sterile glass beads and 40 mL of autoclaved distilled water. The Falcon tubes filled with mycelia materials were then vortex vigorously for 1 min to separate the mycelia into smaller pieces. Mycelial suspension was filtered through 2 layers of cheesecloth to remove the glassbeads and bigger mycelial clumps. The flow-through was then used as mycelial suspension stock (10°) for serial dilution. S-stock of mycelial suspension was further diluted into a series spanning from 10<sup>-2</sup>, 10<sup>-3</sup>, and 10<sup>-4</sup>. This dilution series were plated on PDA using the pour plate method. The number of CFU (colony forming units) was counted and recorded. Mycelial suspensions were adjusted with sterile water to about 10<sup>-5</sup> – 10<sup>-6</sup> CFU/mL for *S. mycoparasitica* and *T. harzianum*, and to about 10<sup>-4</sup> - 10<sup>-5</sup> CFU/mL for *F. graminearum*.

Quantification of interactions between mycoparasite-pathogen-wheat roots was conducted on the spring wheat CDC-TEAL 2001. Wheat plants were grown in containers (4 x 4 x 16 cm) with 10 g of different layers of soil-less growing mix (Figure 10). All the seeds were surface-sterilized prior to sowing. The containers 10 were lined with filter paper and packed with

6 g of Pro-Mix<sup>®</sup> soil-less mix (Sun Gro Horticulture Canada Ltd., Delta, BC, CA; Pro-Mix is a registered trademark of Premier Horticulture Ltd., Riviere-du-Loop, PQ, CA) which comprised the first layer 20. This layer 20 was then overlaid with a second layer 30 (1g) of either Pro-Mix<sup>®</sup> mix amended and homogenized with  $\sim 5$  to  $6 \times 10^4$  CFU of *F. graminearum* mycelial suspension or alternatively with Pro-Mix<sup>®</sup> mix with water only. The second layer 30 was followed by a third layer 40 (1g) of either Sunshine<sup>®</sup> peat moss (Sunshine is a registered trademark of Sun Gro Horticulture Canada Ltd., Seba Beach, AB, CA) supplemented and homogenized with  $\sim 5$  to  $6 \times 10^5$  CFU of *S. mycoparasite* or alternatively with *T. harzianum* mycelial suspension or alternatively with water only. Six spring wheat seeds were then sowed on top of the third layer 40 and topped with a fifth layer 60 of 2 g of Sunshine<sup>®</sup> peat moss (Figure 10). Spring wheat seeds were germinated and grown under a 16-h photoperiod (22°C day/15°C night) with light intensity of  $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ , watered every 2 d, and fertilized every 14 d using 1300 ppm of NPK (20-20-20) fertilizer. All treatments in the experiment were in three replicates and the experiment was repeated twice.

At the mid-seedling growth, corresponding to Zadok's growth stage 13 (Zadoks et al. 1974, Weed Research 14:415-421), wheat plants with their roots were removed from the pots and washed under running tap water to remove all the soil particles. Washed roots were dried with filter papers. Number of germinated seeds, total biomass, root biomass, total length, and root length were counted and measured. Percentage of seed germination was calculated with the following formula: (Number of germinated seeds in particular treatment/number of germinated seeds in control treatment) x 100%. The roots were then subjected to total DNA extraction with DNeasy<sup>®</sup> Plant Mini Kit (Qiagen Inc., Mississauga, ON, CA). Extracted total DNAs from roots of different treatments were employed in real-time PCR quantification.

### *Statistical analyses*

Root biomass (g), total biomass (g), root length (cm), total length (cm), seed germination (%); and *S. mycoparasitica*, *T. harzianum*, and *F. graminearum* genomic DNA quantification from the roots of Spring wheat plants were analyzed by using analysis of variance (ANOVA).  $\text{Log}^{10}$  transformations were carried out whenever required to meet the ANOVA requirements.

Multiple comparisons for more than two samples were analyzed by utilizing Tukey's studentized range test at  $P = 0.05$  (SPSS 1990).

#### *Wheat growth and fungal inoculation*

Root biomass, total biomass, root length, total length, and seed germination of *F. graminearum* infected spring wheat were significantly increased with the treatments of *S. mycoparasitica* compared to inoculation with *F. graminearum* alone (Table 3).

Table 3: Effects of SMCD2220-01 (SM) and *F. graminearum* 3-ADON (Fgra) inoculation treatments on root biomass (g), total biomass (g), root length (cm), total length (cm) and seed germination (%) of spring wheat plants

Treatment	Root biomass (g)	Total biomass (g)	Root length (cm)	Total length (cm)	Seed germination (%)
Control	0.27 ± 0.06 b	0.56 ± 0.56 b	9.63 ± 0.74 b	31.88 ± 2.39 c	NA*
SM	0.32 ± 0.03 ab	0.69 ± 0.69 a	13.88 ± 1.11 a	38.13 ± 2.76 ab	101 ± 2 a
Fgra	0.18 ± 0.03 c	0.38 ± 0.48 c	7.88 ± 1.89 c	23.75 ± 1.98 d	31.25 ± 1.4 c
SM-Fgra	0.27 ± 0.03 b	0.54 ± 0.54 b	12.13 ± 1.35 ab	32.13 ± 3.91 be	87.25 ± 2 b

\* Values expressed are means of six replicates ± standard deviation of the mean. Values followed by same letters within each column are not significantly different using Tukey's range test ( $P < 0.05$ ).

Table 3, the mycoparasite *S. mycoparasitica* demonstrates both biomass stimulation and bioprotection or biocontrol.

#### *Conformation of the Sphaerodes-specific primer set*

The SmyITSF/R primer set was tested with *S. mycoparasitica*, seven *Fusarium* species, eight different ascomycetous fungal isolates two zygomycete fungi, and three basidiomycetous fungal strains. This primer set only amplified *S. mycoparasitica*.

Figure 11 shows SmyITSF/R primers amplified PCR products for *S. mycoparasitica* (SM), five *Fusarium* strains (Fa = *F. avenaceum*, Fo = *F. oxysporum*, Fs = *F. sporotrichioides*, Fg3 = *F. graminearum* chemotype 3, and Fg15 = *F. graminearum* chemotype 15), two *Trichoderma* species (T22 = *T. harzianum* T22 and Tv = *T. viride*), two *Cladosporium* species

(CC = *C. cladosporioides* and CM = *C. minourae*), and *Penicillium aurantiogriseum* (PA) were electrophoresed on 1% agarose gel at 100 V for 20 minutes. The size of the band is around 300 to 400 bp.

#### *Standard curves*

The standard curves based on known diluted concentrations of DNAs from *S. mycoparasitica*, *F. graminearum*, and *T. harzianum* were constructed. Standard curves were achieved using a series of 10-fold diluted DNA spanning from  $3.8 \times 10^2$  to  $3.8 \times 10^{-2}$  ng for *S. mycoparasitica*,  $2.7 \times 10^3$  to  $2.7 \times 10^{-1}$  ng for *F. graminearum* chemotype 3, and  $7.0 \times 10^2$  to  $7.0 \times 10^{-2}$  ng for *T. harzianum*. Quantification demonstrated linear relation ( $r^2 = 0.999$  for *S. mycoparasitica*,  $r^2 = 0.998$  for *F. graminearum*, and  $r^2 = 0.996$  for *T. harzianum*) between  $\log_{10}$  of fungal genomic DNA (in ng/ $\mu$ l) and real-time PCR threshold cycles (Ct) (threshold fluorescence signal of 0.025 was used for all three fungal isolates) (Figures 12A, 12B, and 12C).

Figure 13 shows RT-PCR sigmoidal coloured curves for *Sphaerodes mycoparasitica* (SMCD 2220-01), with 0.025 fluorescence line, showing the range of  $3.8 \times 10^2$  to  $3.8 \times 10^{-2}$  ng in a ten-fold decreasing manner.

#### *RT-PCR confirmation of the Sphaerodes-biocontrol effects*

Real-time PCR, which evaluated quantity of *S. mycoparasitica* SMCD2220-01 and *F. graminearum*, confirmed that amounts of *F. graminearum* DNA detected in treatments with *S. mycoparasitica* SMCD2220-01 and *T. harzianum* were significantly reduced (Figure 14A). Previously, treatments with mycoparasitic *Sphaerodes retispora* had been observed to show significant suppression of *F. oxysporum* in watermelon plants (Harveson et al. 2002, Plant Dis. 86: 1025-1030). The amount of *S. mycoparasitica* DNA detected was not significantly different between wheat inoculated with *F. graminearum* or without *Fusarium* (Fig.14B). The amount of *T. harzianum* DNA detected in the treatment inoculated with *F. graminearum* was observed to be significantly reduced, as compared to non-*Fusarium* treatment (Fig.14C).

In conclusion, during in vitro culture-based studies, only *S. mycoparasitica* SMCD2220-01 was observed to enhance wheat seed germination and formation of secondary roots, whereas T-22 induced post-emergence damping-off symptoms. Under controlled

conditions in pytotron, *S. mycoparasitica* SMCD2220-01 was able to reduce the quantity of *F. graminearum* in spring wheat root, as well as improving the survival and growth of the spring wheat seedlings. In contrast to *T. harzianum*, the amount of *S. mycoparasitica* SMCD2220-01 DNA detected was not significantly different between wheat inoculated with *F. graminearum* and without *Fusarium*. Hence, *S. mycoparasitica* SMCD2220-01 could be a better biocontrol candidate for the *F. graminearum* pathogen in wheat.

#### Example 6

Quantitative RT-PCR results from testing Mycoparasite-*Fusarium* interaction on Wheat host using growing conditions disclosed herein also confirmed the efficiency of *S. mycoparasitica* SMCD2220-01 for significantly decreasing an accumulation of genes implicated in *Fusarium* DON-mycotoxin production.

Strain used: *F. graminearum* 3-ADON SMCD2243, *F. graminearum* 15-ADON SMCD2244 and *S. mycoparasitica* SMCD2220-01 strains.

Primersets used: Tox5-1/2 for *Fusarium* (Wu *et al.* 2002. J. Environ Monit. 4:377-382) and SmyITSF/R (disclosed herein) for *S. mycoparasitica*.

Results are summarized in the Figure 15. Real-time fluorescence curves of *tri5* gene sequences amplified by using Tox5-1/2 primer set from total DNA extracted from dual-culture assays of *Fusarium graminearum* strains and pre-inoculated *Sphaerodes mycoparasitica* SMCD2220-01 (SM) or singly grown *F. graminearum* 3-ADON and 15-ADON chemotypes.

#### Example 7

The effects of filtrates collected from different *Fusarium sp.* on the germination of *Sphaerodes* ascospores were assessed in this study. Also assessed were the ascospore germination patterns.

*Media, Reagents, and chemicals:* Potato dextrose agar (PDA), potato dextrose broth (PDB), yeast extract, malt extract agar (MEA), agar, and peptone were purchased from Difco (Becton



Dickinson Diagnostics, Sparks, Maryland). Streptomycin sulphate, Neomycin sulphate, and other reagents of analytic grade were from Sigma-Aldrich (Oakville, ON, CA).

*Fungal strains and growth conditions:* Four phytopathogenic *Fusarium* strains (*F. avenaceum* SMCD 2241, *F. oxysporum* SMCD 2242, *F. proliferatum* (Matsush.) Nirenberg SMCD 2244, and *F. sporotrichioides* Sherb SMCD 2243), three beneficial fungal inoculants: *T. harzianum* (RootShield<sup>®</sup>), *P. bilaii* ([JumpStart<sup>®</sup>), and *Chaetomium globosum* Kunze; and one mycoparasitic fungal strain, *S. mycoparasitica* were maintained on PDA amended with antibiotics (100 mg/L streptomycin sulphate and 12 mg/L neomycin sulphate) and used throughout this study. An isolate of the mycoparasitic fungus, *S. mycoparasitica*, was separated from its *F. oxysporum* host on myclobutanil agar (MBA) selective medium as previously described and a monosporic culture was achieved by picking up mature perithecia which were then suspended and shaken in 2-mL sterile distilled water blanks to encourage release of ascospores. The ascospore-suspension was then spread on PDA plates. After a few minutes, a Carl Zeiss Stemi 1000 dissecting microscope was used to identify individual ascospores, which were removed and transferred to PDA supplemented with 100 mL/L of *Fusarium* filtrate (*F. avenaceum* and *F. oxysporum*). All fungal isolates were maintained in the culture collection of the Saskatchewan Microbial Collection and Database, Canada (SMCD).

*Ascospore production:* Mycelium plugs from the margin of the actively growing monosporic-derived *S. mycoparasitica* culture were cut and placed on Modified Leonian's Agar (MLA: maltose, 6.25g; malt extract, 6.25g; KH<sub>2</sub>PO<sub>4</sub>, 1.25g; yeast extract, 1.0g; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.625g; peptone, 0.625g; agar, 20g, and 1L of dH<sub>2</sub>O) (Malloch, D., and Cain, R.F. (1971) New genera of Onygenaceae. Canadian Journal of Botany 49: 839-846). Inoculated Modified Leonian's Agar (MLA) plates were incubated at room temperature (23 °C) under darkness conditions for a month before collecting spores for germination assays. Mature ascospores were collected according Goh, Y.K. and Vujanovic, V. ((2010) *Sphaerodes quadrangularis* biotrophic mycoparasitism on *Fusarium avenaceum*. Mycologia 102(4):757-62).

*Preparation of the fungal filtrates:* Four pathogenic *Fusarium* strains and three beneficial fungal strains were grown in shake cultures for 7 d at room temperature (23 °C) (with 250 rpm) in 500 mL flasks, each with 100 mL PDB medium prior to fungal filtrates extraction. After incubation, the mycelia were removed by filtering through filter paper (Whatman<sup>®</sup> Grade No. 1) and the filtrates were then filter-sterilized with 0.02-µm-pore-size nitrocellulose filter (Fisher Scientific

Ltd., Nepean, ON, CA). Only fresh preparations of fungal filtrates were employed for the spore germination test in this study.

*Effect of heat and cold treatments on S. mycoparasitica spore germination:* Heat activation and cold-shock treatments were performed to investigate ascospore germination in *S. mycoparasitica*. Aqueous ascospore suspensions were heat-shocked at 60 and 65 °C for 20 min, and cold-treated at 4, -20, and -70 °C for 5 min and 20 min. The heat- and cold-shocked spores (10 µL) were then transferred and inoculated onto water agar (WA) and PDA for 1 d and 3 d. Spore germination was checked daily. Spores not subjected to heat and cold treatments were used as controls. All treatments were in three replicates and experiment was repeated twice.

No germination was observed in all the heat and cold treatments over a 7 d period.

*Germination on various media:* Aqueous spore suspensions (10 µL) of *S. mycoparasitica* were transferred and inoculated onto the surface of the following media: 1.5% water agar (WA), PDA, MLA, MEA, Carnation leaf agar (CLA) (Tschanz et al. 1976), 1.5% water agar plus 100 mL/L of *Fusarium* filtrate, and PDA with 100 mL/L of the *Fusarium* filtrate. *Fusarium* filtrates were created using *F. avenaceum*, *F. oxysporum*, *F. proliferatum*, and *F. sporotrichioides*. *S. mycoparasitica* ascospore-inoculated plates were then incubated at room temperature (23 °C) for 3 d, and spore germination was examined daily. All treatments were in three replicates and experiment was repeated twice.

There was no spore germination on 1.5% WA, PDA, MLA, MEA, and CLA. The effects of 1.5% WA and PDA media amended with 100 mL/L of *Fusarium*-filtrates on spore germination were examined and are summarized in Table 4.

No germination was recorded on day 3 initially and observations were continued up to 7 d on WA or PDA alone and WA or PDA with either *F. proliferatum* or *F. sporotrichioides*. However, when either *F. avenaceum* or *F. oxysporum* filtrates were added to WA and PDA, the percentage of ascospore germination in *S. mycoparasitica* drastically increased after 3 d incubation. On day 3, germination increased on the PDA amended with *Fusarium*-filtrate, compared to growth on WA supplemented with *Fusarium*-filtrate, for *S. mycoparasitica*

Table 4: Percentage germination of *Sphaerodes mycoparasitica* ascospore on various types of *Fusarium*-filtrate supplemented media, including water agar and potato dextrose agar checks

<i>S. mycoparasitica</i> Medium	Spore germination (%)*	
	Day 1**	Day 3**
Water agar (WA)	0 <sup>d</sup>	0 <sup>d</sup>
<i>F. avenaceum</i> -WA	11.1 ± 1.7 <sup>b</sup>	35.2 ± 2.6 <sup>b</sup>
<i>F. oxysporum</i> -WA	1.6 ± 1.3 <sup>c</sup>	18 ± 1.4 <sup>c</sup>
<i>F. proliferatum</i> -WA	0 <sup>d</sup>	0 <sup>d</sup>
<i>F. sporotrichioides</i> -WA	0 <sup>d</sup>	0 <sup>d</sup>
Potato dextrose agar (PDA)	0 <sup>d</sup>	0 <sup>d</sup>
<i>F. avenaceum</i> -PDA	21.5 ± 1.4 <sup>a</sup>	63.2 ± 2.4 <sup>a</sup>
<i>F. oxysporum</i> -PDA	2.5 ± 2 <sup>c</sup>	37.5 ± 2 <sup>b</sup>
<i>F. proliferatum</i> -PDA	0 <sup>d</sup>	0 <sup>d</sup>
<i>F. sporotrichioides</i> -PDA	0 <sup>d</sup>	0 <sup>d</sup>

\* Numbers in each column represented mean of ascospore germination (in %) ± standard deviation.

\*\* Each incubation day for *Sphaerodes* was analyzed separately. Means within each column for each medium treatment followed by the same letter are not significantly different at  $P \leq 0.05$  after Kruskal-Wallis test.

*Effects of fungal filtrates on spore germination:* Fungal filtrates from four different phytopathogenic *Fusarium* species and three beneficial fungal isolates were employed to study spore germination of *S. mycoparasitica*, and host specificity response. Aqueous spore suspensions of *S. mycoparasitica* were suspended in the filtrates of four separate pathogenic *Fusarium* spp. and three beneficial fungal strains for 1 d and 3 d in the ratio of 1:2 (1 part of aqueous spore suspension: 2 parts of fungal filtrate). Filtrate-suspended spores (10 µL) were then transferred and inoculated onto PDA for an additional day (spore germination was counted on PDA plate). Spore germination was count according to Goh and Vujanovic (2010) after an additional day of inoculation on PDA medium. Control treatments were suspended with sterilized distilled water or PDB. Abbreviations for different treatments at four separate chronosequences throughout the experiment: 1d Sus, 1d Sus + 1d PDA, 3d Sus, and 3d Sus + 1d PDA represent 1 day *Fusarium*-filtrate suspension, 1 day filtrate suspension with an additional day incubation on PDA medium, 3 day suspension, and 3 day filtrate suspension with an additional day incubation on PDA, respectively.

No spore germination was observed in the treatments with *P. bilaii* and *C. globosum* fungal filtrates. Both water and PDB suspension controls appeared to trigger approximately 1.8-3.8% spore germination for *S. mycoparasitica*, on day 3 with an additional day on PDA only (Table 5). Ascospores of *S. mycoparasitica* suspended in filtrates of pathogenic *F. sporotrichioides* and beneficial *T. harzianum* showed no germination on day 1 in suspension, but low number of ascospore germination was observed after 1 day in suspension followed by an additional day on PDA, and in the other incubation treatments. Ascospores suspended in *F. proliferatum*-filtrate were observed to germinate in low abundance for 1 d, 1 d with an additional day on PDA, 3 d, and 3 d plus an additional day on PDA. *Sphaerodes* ascospores showed the highest number of germinated ascospores in the *F. avenaceum*-filtrate suspension. In the day 1 suspension treatments, ascospore germination of *S. mycoparasitica* in *F. avenaceum*-filtrate was significantly higher (89.2%) than in other treatments. When ascospores of the *S. mycoparasitica* were suspended in *F. oxysporum*-filtrate, the amount of spore germination was increased compared to control. There was no germination recorded for the 1 d suspension in the *F. oxysporum*-filtrate treatment. However, a low number of ascospores were stimulated in *F. oxysporum*-filtrate suspension on the second day. Based on the ascospore germination rate and response, *S. mycoparasitica* showed high specificity to *F. avenaceum* and *F. oxysporum*.

Table 5: Percentage spore germination of *Sphaerodes mycoparasitica* ascospores suspended in different *Fusarium* and biocontrol fungi filtrates assessed throughout four chronosequences. Control treatments were suspended with sterilized distilled water (SDW) or potato dextrose broth (PDB).

<i>S. mycoparasitica</i> Treatment**	Spore germination (%)*				
	1d Suspension	1d Sus + 1d PDA	3d Suspension	3d Sus + 1d PDA	3d Sus + 1d PDA
SDW	0 <sup>c</sup>	0 <sup>c</sup>	0 <sup>e</sup>	0 <sup>e</sup>	2 ± 1.2 <sup>c</sup>
PDB	0 <sup>c</sup>	0 <sup>c</sup>	0 <sup>e</sup>	0 <sup>e</sup>	2.8 ± 1.9 <sup>c</sup>
<i>F. avenaceum</i> -filtrate	89.2 ± 6.2 <sup>a</sup>	91 ± 8.4 <sup>a</sup>	93.2 ± 6.7 <sup>a</sup>	94.2 ± 7.9 <sup>a</sup>	94.2 ± 7.9 <sup>a</sup>
<i>F. oxysporum</i> -filtrate	0 <sup>c</sup>	81.8 ± 11 <sup>b</sup>	85 ± 7.9 <sup>b</sup>	91 ± 10 <sup>a</sup>	91 ± 10 <sup>a</sup>
<i>F. proliferatum</i> -filtrate	2.6 ± 3.6 <sup>b</sup>	3.2 ± 4.6 <sup>d</sup>	10 ± 7.4 <sup>c</sup>	11.2 ± 8 <sup>b</sup>	11.2 ± 8 <sup>b</sup>
<i>F. sporotrichioides</i> -filtrate	0 <sup>c</sup>	3 ± 2.8 <sup>d</sup>	3.4 ± 3.3 <sup>d</sup>	10 ± 5.6 <sup>b</sup>	10 ± 5.6 <sup>b</sup>
<i>P. bilaii</i> -filtrate	0 <sup>c</sup>	0 <sup>e</sup>	0 <sup>e</sup>	0 <sup>c</sup>	0 <sup>c</sup>
<i>T. harzianum</i> -filtrate	0 <sup>c</sup>	9 ± 1.1 <sup>c</sup>	1.6 ± 1.6 <sup>d</sup>	10.4 ± 1.5 <sup>b</sup>	10.4 ± 1.5 <sup>b</sup>
<i>C. globosum</i> -filtrate	0 <sup>c</sup>	0 <sup>e</sup>	0 <sup>e</sup>	0 <sup>c</sup>	0 <sup>c</sup>

\* Numbers in each column represented mean of ascospore germination (in %) ± standard deviation.

\*\* Each incubation day for *S. mycoparasitica* was analyzed separately. Means within each column of *S. mycoparasitica* for each filtrate-suspension treatment followed by the same letter (in superscript) are not significantly different at  $P \leq 0.05$  after Kruskal-Wallis test.

*Spore germination assessments:* Microscopic assessments of ascospore germination were conducted after incubation for 1 d and 3 d (spore germination was counted on suspensions). Percentage of germinated spores was obtained by scoring the spores on a Petri dish while observing them through the 200x and 400x objectives of the Carl Zeiss Axioskop2 microscope and systematically choosing 50 spores, starting at the top right corner and continuing to count until 50. Each drop of the ascospore suspension on a growth medium plate was considered as a subunit, and there were three subunits or replicates per plate. The experiments were repeated twice. An ascospore was only considered germinated when the germ tubes exceeded the width of the ascospore (approximately 12 $\mu$ m). Germinated ascospores were counted and recorded as a percentage of the total ascospore number that was counted.

Ascospores from *S. mycoparasitica* showed two kinds of germination patterns. Single polar germination was more prevalent in the *F. oxysporum*-filtrate suspension (Figures 16A, 16B, 16D, 16E). A small number of spores from *S. mycoparasitica* found to produce shorter double-polar germination in the treatment with *F. oxysporum*-filtrate at 3 d suspension with additional 1 d on PDA (Figure 16F). In the *F. avenaceum*-filtrate suspension, ascospores of *S. mycoparasitica* showed higher preference for two-polar germination (Figures 16G). Single polar germination was also found in *F. avenaceum*-filtrate suspension; however it was lower than in *F. oxysporum*-filtrate treatment. Commonly, these single-polar germinated spores produced larger web-like organizations and longer hyphal formation (in *F. avenaceum*-filtrate suspension), which was rarely found in *F. oxysporum*-filtrate suspended spores (Figures 16H, 16I). The majority of the germinated ascospores in *S. mycoparasitica*, which was activated through suspension in *F. oxysporum*-filtrate were detected to be either at an angle of 90° (Figures 16E, 16F) or between 90° and 180° (Figure 16F) at the tip of the polar germ pores. However, very few spores germinated at an angle of 180° (Figure 16D). Most activated spores (with *F. avenaceum*-filtrate treatment) showed germination at an angle of 180° (Figure 16G).

#### Example 8

The host specificity of *S. mycoparasitica* to *F. graminearum* 3-ADON and 15-ADON strains were assessed in this study.

*Media, reagents and chemicals:* Potato dextrose agar (PDA), potato dextrose broth (PDB), yeast extract, malt extract agar (MEA), agar, and peptone were purchased from Difco (Becton Dickinson Diagnostics). Streptomycin sulphate, Neomycin sulphate, and other reagents of analytical grade were from Sigma-Aldrich.(Oakville, ON, CA) IQ SYBR Green Supermix™ for real-time PCR reactions was acquired from Bio-Rad Laboratories (Mississauga, ON, CA).

*Fungal strains and growth:* All phytopathogenic *Fusarium* strains: *Fusarium graminearum* 3-ADON (Fgra3) SMCD 2243, and 15-ADON (Fgra15) SMCD 2244 chemotypes, *F. avenaceum* (Fave) SMCD 2241, *F. oxysporum* (Foxy) SMCD 2242, *F. proliferatum* (Fpro) SMCD 2244, *F. sporotrichioides* (Fspo) SMCD 224; and one mycoparasitic *Sphaerodes mycoparasitica* SMCD 2220 strain were retrieved from Saskatchewan Collection and Database (SMCD), maintained on PDA amended with antibiotics (100 mg/L streptomycin sulphate and 12 mg/L neomycin sulphate) and used throughout this study.

*Spore production and germination assays:* Ascospores of *S. mycoparasitica* were produced on Modified Leonian's agar (MLA), harvested and prepared as previously described. Also, *Fusarium* spp. filtrates were prepared, *S. mycoparasitica* spore germination assays in six different *Fusarium* filtrates were carried out as previously described.

*Sphaerodes mycoparasitica* spore germination suspended in both *F. graminearum* chemotype 3-ADON and 15-ADON filtrates was lower compared to *F. avenaceum* for the first incubation day, and compared to both *F. avenaceum* and *F. oxysporum* for the remaining incubation days ( $P \leq 0.05$ ; with Mann-Whitney Test) (Figure 17). No significant differences in germination between *F. graminearum*, *F. proliferatum*, and *F. sporotrichioides* filtrate treatments were observed for the first two incubation days. However, treatments with *F. graminearum* filtrates showed significantly higher germination rate of *S. mycoparasitica* compared to *F. sporotrichioides* filtrate treatment during later incubation time points (Figure 18).

*Dual-culture assays:* Dual-culture assays for examining the degree of hyphal reduction/inhibition or damage to *F. graminearum* chemotypes were assessed as disclosed in Goh and Vujanovic (2009, Mycologia DOI: 10.3852/69-171). *S. mycoparasitica* is slow-growing fungus as

compared to *F. graminearum* 3 and 15 strains. Therefore, *S. mycoparasitica* was pre-inoculated onto the PDA plates for 1 d, at 21 °C in darkness, prior to inoculating *Fusarium* mycelial plugs. Linear mycelial growth of *Fusarium* strains for both treatments indicated above was measured and recorded daily for 5 days. Around 0.5 x 1.5 cm<sup>2</sup> (sampling zones) located approximately 0.2 cm behind the contact zone between *F. graminearum* and *S. mycoparasitica* was excised and subjected to DNA extraction. Each treatment was with three replicates and the experiment was repeated twice. The PDA plate inoculated with *F. graminearum* only was the positive control. Total genomic DNA was extracted using a DNeasy<sup>®</sup> Plant Mini Kit. The DNA was eluted once in 50 µl of buffer AE and stored at -20°C until real-time PCR quantification assays (as described below).

Since *S. mycoparasitica* demonstrated slower mycelial growth (0.56 cm per day; n = 9) compared to *F. graminearum* 3-ADON (0.74 cm per day; n=6) and 15-ADON (0.68 cm per day; n=6) chemotypes, the linear growth of *F. graminearum* mycelia in dual-culture was assessed using the pre-inoculation method. *S. mycoparasitica* was pre-inoculated on PDA for 1 d followed by *F. graminearum* inoculation. The pre-inoculation approach demonstrated significant differences (starting day 3) in linear growth suppression of *F. graminearum* chemotype 3 and 15 compared to the co-inoculation approach (Figures 19A and 19B).

*Establishment of mycoparasitism:* Fusion biotrophic mycoparasitic interactions between *S. mycoparasitica* and both *F. graminearum* chemotype strains, and intracellular parasitism interactions were examined and assessed on slide cultures according the methods described in Goh & Vujanovic (2009).

On day 3 of inoculation on PDA with *F. graminearum* 3-ADON and 15-ADON, no clamp-like or hook-like structures were formed by *Sphaerodes mycoparasitica* on *Fusarium* strains. On day 5 of inoculation, clamp-like and hook-like contact structures as well as *Fusarium* hyphal cells penetration (with haustoria) were observed (Figures 20E to 20I). Furthermore, on day 3, *S. mycoparasitica* removed red pigment from the mycelia of *F. graminearum* 3-ADON on the slide culture (Figures 20A to 20D). As a result, *S. mycoparasitica* mycelia adopted reddish color (Figure 20C). Between day 4 and 5, formation of red crystal-like pellets was detected on the surface of mycoparasite hyphae (Figure 20D). The mechanism behind the color changes



observed remains unknown. For *F. graminearum* chemotype 15-ADON, no uptake of red complex or release of red crystal-like structures by *S. mycoparasitica* hyphae were noted. Nevertheless, flower-like hyphal structures appeared which could indicate possible growth inhibition of 15-ADON *F. graminearum* (Figure 20J). Significant differences in diameters of infected and non-infected hyphae were seen for both *F. graminearum* chemotypes (Figure 21).

*Primers and standard curves:* *F. graminearum*-specific (Fg16NF/R) and trichothecene *Tri5* gene-specific (Tox5-1/2) primer sets were used in this study. Standard curves for *F. graminearum*- and *Tri5* gene-primer sets were generated, based on threshold cycles (Ct), by using a series of 10-fold diluted genomic DNAs from *F. graminearum* (spanning from  $2.7 \times 10^2$  to  $2.7 \times 10^{-2}$  ng/ $\mu$ l for *F. graminearum*-specific primer set, from  $2.7 \times 10^2$  to  $2.7 \times 10^{-1}$  ng/ $\mu$ l of 3-ADON strain and  $3.0 \times 10^1$  to  $3.0 \times 10^{-2}$  ng/ $\mu$ l of 15-ADON strain DNAs for *Tox5-1/2* primer set). Ct values were recorded and obtained by the Opticon Monitor™ software version 3.1 (Bio-Rad Laboratories Inc., Mississauga, ON, CA). Standard curves for different primer sets were constructed by plotting the threshold cycles (Ct) value versus the logarithm ( $\log_{10}$ ) of the concentration of 10-fold serial diluted *F. graminearum* DNAs as described above. Amplifications with different primer sets on the genomic DNAs of two *F. graminearum* chemotypes were run in triplicates to obtain the mean and standard deviation of each 10-fold serial dilution.

*Real-time PCR quantification:* Real-time PCR amplifications on total genomic DNA extracted from the sampling zones (as described above) were performed using MiniOpticon™ (Bio-Rad Laboratories Inc., Mississauga, ON, CA). All the real-time PCR reactions were performed by utilizing the real-time PCR MJ white tubes (Bio-Rad Laboratories Inc., Mississauga, ON, CA) in a total volume of 25  $\mu$ l. The reaction mixture for all real-time PCR assays were: 12.5  $\mu$ l of IQ Supermix (Bio-Rad Laboratories Inc., Mississauga, ON, CA), 1  $\mu$ l of each 10  $\mu$ M forward/reverse primers (Invitrogen), 9.5  $\mu$ l of sterilized UltraPure Millipore water, and 1  $\mu$ l of DNA template. Real-time PCR conditions for Fg16NF/R primer set used were outlined in Nicholson *et al.* (1998, *Physiol. Mol. Plant Pathol.* 53: 17-37) with melting curve analysis at 60 to 95°C. Parameters for Tox5-1/2 primer set as described in Schnerr *et al.* (2001, *Int. J. Food microbiol.* 71: 53-61).

Standard curves for different primer sets with different *F. graminearum* DNA sources were constructed (Figure 22). Growth suppression or inhibition at the sampling zones (Figure 23) for *F. graminearum* chemotype 3 and 15 was further confirmed with real-time PCR amplifications with *F. graminearum*- and *tri5* gene-specific primer sets. Sigmoidal curves for the four different treatments (*F. graminearum* chemotype 3 or 15 only and *F. graminearum* chemotype 3 or 15 pre-inoculated with *S. mycoparasitica*) with Fg16NF/R primer set were generated using Opticon Monitor software version 3.1 and illustrated in Figure 24.

Using Fg16NF/R primer set, quantity of *F. graminearum* chemotype 3 DNA in the sampling zones significantly decreased when pre-inoculated with *S. mycoparasitica* compared to un-inoculated treatment ( $P = 0.01$ ) (Figure 25). However, DNA of *F. graminearum* chemotype 15 did considerably but not significantly reduced ( $P = 0.085$  using T-test). Using Tox5-1/2 primer set, amount of *tri5* gene fragments diminished appreciably in both *F. graminearum* chemotype 3 and 15 challenged with *S. mycoparasitica* ( $P \leq 0.05$ ).

#### Example 9

The effects of *S. mycoparasitica* (biotrophic mycoparasite) and *T. harzianum* (necrotrophic mycoparasite) on expression of *Tri* and *PKS* genes by *F. graminearum* 3-ADON and 15-ADON strains were assessed in this study.

*Fungal strains and growth:* Two *Fusarium graminearum* 3-ADON (SMCD 2243) and 15-ADON (SMCD 2244) chemotypes, *Trichoderma harzianum* necrotrophic (SMCD 2166) and *Sphaerodes mycoparasitica* biotrophic (SMCD 2220) mycoparasites were obtained from the Saskatchewan Microbial Collection and Database (SMCD). All strains were maintained on Potato dextrose agar (PDA, BD Biosciences, Mississauga, ON, CA) amended with antibiotics (100 mg/L streptomycin sulphate and 12 mg/L neomycin sulphate; Sigma-Aldrich Canada Ltd., Oakville, ON, CA) prior to study initiation.

*Chemical fungicide control:* A concentration of 100  $\mu\text{mol/L}$  tebuconazole was prepared from Folicur<sup>®</sup> 432F (43.2% tebuconazole, Bayer CropScience Inc., Saskatoon, SK, CA; Folicur is a

registered trademark of Bayer Aktiengesellschaft, Leverkusen, Fed. Rep. Germany). This fungicide preparation was used throughout the study.

*In vitro* assay, sampling and RNA extraction: Dual-culture assay was carried out between *F. graminearum* and Folicur<sup>®</sup> (100 µmol/L tebuconazole) or biological (*T. harzianum* or *S. mycoparasitica*) agents on Minimal medium as disclosed by Xue et al. (2009, Can. J. Plant Pathol. 31: 169-179). Inoculated dual-culture plates were incubated at room temperature (23C) in dark conditions for a week. Mycelia of *F. graminearum* were harvested after a week of inoculation with either chemical or biological agents. A 0.5-cm<sup>2</sup> mycelial plug was cut out approximately 0.5 cm away from the border of interaction and fungal cells were disrupted with liquid nitrogen. Total RNA from the samples were extracted using Aurum Total RNA mini kit<sup>™</sup> and extracted RNA was treated with DNase (Bio-Rad Laboratories Inc., Mississauga, ON, CA) according the manufacturer's recommendations. Samples were then stored at -70C till gene expression analysis.

During the *In vitro* studies, both *F. graminearum* 3-ADON and 15-ADON chemotypes changed mycelia morphology corresponding to each treatment (Figures 26A - 26B). However, the main diagnostic distinction was the tendency of *F. graminearum* chemotypes to abundantly produce chlamidospores in clusters or chains when exposed to tebuconazole fungicide compared to mycoparasites (Figures 26C - 26D). In the case of *F. graminearum* chemotype 15 (15-ADON producer), all four trichothecene genes – *Tri4*, *Tri5*, *Tri6*, and *Tri10* in this *Fusarium* strain were found to be induced in high amounts when treated with chemical tebuconazole compared to treatments that co-inoculated with biological agents (either *S. mycoparasitica* or *T. harzianum*) (Figures 27A - 27D). *F. graminearum* chemotype 3 (3-ADON producer) was observed to demonstrate induction of *Tri4*, *Tri5* and *Tri10* genes when co-inoculated with necrotrophic mycoparasitic *Trichoderma harzianum* (Figures 27A, 27B, 27D). When *F. graminearum* chemotype 3 was treated with Folicur<sup>®</sup> fungicide, only *Tri10* gene was being induced (Figure 27D). Generally, all four *Tri* genes were repressed significantly when challenged with biotrophic mycoparasitic *Sphaerodes mycoparasitica* (Figure 27A - 27D).

*Real-time Reverse-transcription PCR*: Real-time RT-PCR was performed by using an IScript One-Step RT-PCR kit<sup>™</sup> with SYBR Green on a MiniOpticon Cyclor System (Bio-Rad

Laboratories Inc., Mississauga, ON, CA), according to the manufacturer's instruction. Primer sets used for amplification and gene expression were summarized in Table 6.

Table 6: Primer sets used to amplify *Tri4*, *Tri5*, *Tri6*, *Tri10*, *PKS4*, *PKS13*, and  $\beta$ -*tubulin* by Real-time PCR.

SEQ ID	Gene	Primer sequences	Reference
SEQ ID NO: 6/7	<i>Tri4</i>	Tri4-F: TAAACGCCCGCGAAGTTCACA Tri4-R: TGGTGATGGTTCGCTTCGAG	Jiao et al. 2008, FEMS Lett. 285: 212-219
SEQ ID NO: 8/9	<i>Tri5</i>	Tr5F: AGCGACTACAGGCTTCCCTC Tr5R: AAACCATCCAGTTCTCCATCTG	Doohan et al. 1999, Appl. Environ. Microbiol. 65: 3850-3854
SEQ ID NO: 10/11	<i>Tri6</i>	Tri6-1 TCTCTACCAACGGTGGATTCAACC Tri6-2 AGCCTTTGGTGCCGACTTCTTG	Pinson-Gadais et al. 2008, Mycopathol. 165: 51-59.
SEQ ID NO: 12/13	<i>Tri10</i>	Tri10-F: TCTGAACAGGCGATGGTATGGA Tri10-R: CTGCGGCGAGTGAGTTTGACA	Jiao et al. 2008
SEQ ID NO: 14/15	<i>PKS4</i>	PKS4-PS.1 GTGGGCTTCGCTAGACCGTGAGTT PKS4-PS.2 ATGCCCTGATGAAGAGTTTGAT	Lysøe et al. 2006, Appl. Environ. Microbiol. 72:3924-3932
SEQ ID NO: 16/17	<i>PKS13</i>	PKS13-PS.1 CCCCCAACTCGACGTCAAATCTAT PKS13-PS.2 TTCTTCCCGCCGACTTCAAACA	Lysøe et al. 2006
SEQ ID NO: 18/19	$\beta$ - <i>tubulin</i>	FGtubf GGTCTCGACAGCAATGGTGTT FGtubr GCTTGTGTTTTTCGTGGCAGT	Lysøe et al. 2006

RT-PCR sample (~25  $\mu$ L) contained 3  $\mu$ L of RNA template, 8.85  $\mu$ L of nucleae-free water, 12.5  $\mu$ L of RT-PCR reaction mixture (2x), 0.5  $\mu$ L of IScript RT enzyme mix (50x) (Bio-Rad Laboratories Inc., Mississauga, ON, CA), 0.1  $\mu$ L of 50  $\mu$ M solutions of both forward and reverse targeted gene-specific primers (Invitrogen Corp., Carlsbad, CA, USA). Real-time PCR conditions were performed as outlined by manufacturer's recommendations: 50C for 10min, 95C for 5min, followed by 40 cycles of denaturing at 95C for 10s and annealing at 55C for 30s, 95C for 1min and 55C for 1 min. PCR reactions were checked for absence of any primer-dimer formation or non-specific PCR amplification by performing melting curve analysis. Contamination of RNA template with residual genomic DNA was eliminated because there was no amplification detected using reverse transcriptase free real-time RT-PCR reaction as template. Fold change in gene expression for each treatment was normalized to  $\beta$ -*tubulin* internal reference gene and relative to the expression for the control treatment (*Fusarium* alone on minimal medium), using the  $2^{-\Delta\Delta CT}$  method proposed by Livak, K.J., and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  Method. Methods 25(4):402-408.  $\Delta\Delta C_T = (C_{T, target}$

gene -  $C_{T, Fgtub}$ )<sub>treatment</sub> - ( $C_{T, target-gene} - C_{T, Fgtub}$ )<sub>control</sub>, where treatment and control indicate *F. graminearum* challenged with chemical or biological agent and *F. graminearum* alone, respectively.

For gene expression analyses based on targeted genes (*PKS4* and *PKS13*) that responsible for zearalenone biosynthesis, these two polyketide synthase genes from both *F. graminearum* chemotypes were detected to be repressed in all chemical or biological treatments (Figures 28A - 28B). Repression of *F. graminearum* chemotype 15 *PKS4* and *PKS13* genes in the treatment with *S. mycoparasitica* biotrophic mycoparasite was illustrated to be significantly higher compared to treatments with *T. harzianum* and tebuconazole fungicide (Figures 28A - 28B).

*Tri* genes were observed to be more sensitive in *F. graminearum* chemotype 3 when co-inoculated with *Trichoderma* necrotrophic mycoparasite, however, in *F. graminearum* chemotype 15, *Tri* genes appeared to be more responsive towards treatment with chemical fungicide (Figures 27A-27D).

*PKS4* gene for both *F. graminearum* chemotypes was monitored to be more sensitive towards treatments with *Trichoderma* necrotrophic mycoparasitic fungus compared to chemical and biotrophic mycoparasitic agents (Figure 28A). However, *PKS13* gene in *F. graminearum* chemotype 15 was found to demonstrate higher sensitivity towards chemical stimulus (Figure 28B).

*Mycotoxins extraction and analyses:* DON, ZEA, 3-ADON and 15-ADON mycotoxins were extracted from agar mycelial plugs (0.5cm<sup>2</sup>) cut from the sampling zone located approximately 0.5 cm behind the contact zone between *F. graminearum* and *S. mycoparasitica*. The extraction was performed by three 10-ml volumes of ethyl acetate. The samples were sonicated on ice and shaken vigorously in ethyl acetate, and then they were allowed to stand for 5 min for separation of phases. The organic phase was siphoned off and passed through sodium sulfate to remove water. The solvent was allowed to evaporate at room temperature (23 °C) for 3 days. The residue was then re-dissolved in 2 ml of acetonitrile for thin liquid chromatography (TLC) (only ZEA) and high performance liquid chromatography (HPLC) (all mycotoxins) analyses. TLC was performed on Silica gel 60 plates (Merck Frosst Canada Ltd, Kirkland, PQ, CA) in the solvent

system with ethyl acetate-toluene (3:1) for 40 min. 3-ADON on the gel plates was sprayed with 20% aluminum chloride in 95% ethanol, detected and analyzed following the protocols disclosed by Vasavada and Hsieh (1987, Appl. Microbiol. Biotech. 26: 517-521). Known standard controls of pure DON, ZEA, 3-ADON and 15-ADON (Sigma-Aldrich Canada Ltd., Oakville, ON, CA) were employed to compare with other treatments. All four mycotoxins were quantified using a Waters' 2695 HPLC system with: 250 x 4.60 mm, Luna 5 $\mu$  micron C18 (2) 100A column (Phenomenex Inc., Torrance, CA USA) and a photodiode-array (PDA) detector was used with an isocratic solvent system (methanol: water-methano containing 5% (v/v) (90:10) ratio). The PDA detector measured the UV spectrum (190-500 nm). Samples were dissolved in acetonitrile and 10  $\mu$ l loaded onto the column using automatic injector and mycotoxins were eluted with solvent as a mobile phase at a rate of 0.75 ml min<sup>-1</sup> for 25 mins. Peak height method was incorporated to determine the exact amount of DON, ZEA, 3-ADON and 15-ADON against a standard curves. Ratio between mycotoxins extracted from treatments of *F. graminearum* treated with Folicur and *F. graminearum* only was calculated for HPLC.

The amount of DON produced was between 70 to 90 ug/L. Consequently, it was too low to be detected in TLC analysis. ZEA was high enough to be analyzed with TLC and both *F. graminearum* chemotypes were observed to produce higher amounts of ZEA toxin under the treatment with Folicur<sup>®</sup> compared to other treatments (Figure 29). With HPLC, ZEA was found to be reduced when challenged with Folicur<sup>®</sup> compared to *F. graminearum* colony only (Figure 30). DON and 3-ADON was detected to increase for both *F. graminearum* when inoculated together with Folicur<sup>®</sup> (Figure 30). Folicur<sup>®</sup> was monitored to trigger highest amount of 15-ADON under *in vitro* assay (Figure 30).

#### Example 10

The effects of *S. mycoparasitica* (biotrophic mycoparasite) and *T. harzianum* (necrotrophic mycoparasite) on expression of aurofusarin gene by *F. graminearum* 3-ADON and 15-ADON strains were assessed in this study.

*Fungi, media, and culture conditions:* Fungal strains were obtained from culture collections at University of Saskatchewan (Food and Bioproduct Sciences fungal collection). Strains of the following fungal species were used, plant pathogenic *Fusarium graminearum*, *F. graminearum*

(3ADON), *F. graminearum* (15ADON), *F. avenaceum*, *F. culmorum*, , *F. proliferatum*, *F. oxysporum*, *F. arthrosporoides* and mycoparasitic *Spaerherodes mycoparasitica* and *Trichoderma harzianum*. Fungi were maintained on potato dextrose agar for 2 wks at 21°C in darkness. Folicur<sup>®</sup> was also used in this study.

Experiments for the impact of red color nuances on aurofusarin gene expression was carried out using previously classified *F. avenaceum* isolates, based on VCGs and colony colours. *F. avenaceum* isolates color identification numbers (CIN) were generated with Hex Color Code Chart. CINs are: Ds #71232B: Red and highly virulent, tolerant at 80°C for 4 hours; Es #4E040B: Moderately red and moderately virulent, tolerant 40°C for 4 hours; Bs #A86608: White and non virulent, susceptible 40°C for 4 hours. Three 0.1 g mycelium samples from three replicates of each isolate was mixed and used for DNA extraction and RT-PCR analyses.

*Fungal DNA Extraction and PCR:* Fungal cultures were grown on 1.5 ml PD broth and centrifuged at 10,000 rpm for 5 min. Supernatant was discarded and, subsequently a total DNA of 2-week old cultures representative of each phenotype/VCG was extracted from the pellet with an Ultra Clean microbial DNA Isolation Kit™ (Qiagen Inc., Mississauga, ON, CA) following manufacture's instructions. The purified DNA was resuspended in 50 µl of elution buffer and stored at -20° C until further analyses. Full length genomic sequences of PKS region (7.2Kb) from *Fusarium graminearum* were obtained from NCBI Genbank. Six pairs of primers were designed (Table 7).

Table 7: PCR Primers designed from full-length genomic sequences of the PKS region of *F. graminearum*

SEQ ID	Primer	Sequence
SEQ ID NO: 20	PKSF1	TCGAGTTTCGTGTTGCGTGT
SEQ IS NO. 21	PKSR1	AGGTAGTTCGCCATACCCGT
SEQ ID NO: 22	PKSF2	AATTGTGCCCGAGGCAGTAC
SEQ ID NO: 23	PKSR2	CATTGGTTCCGCCCGCAATAG
SEQ ID NO: 24	PKSF3	TGACAACTTCGCTGGTTTGGG
SEQ ID NO: 25	PKSR3	CATAGCTTGGCCAGTGCCATC
SEQ ID NO: 26	PKSF4	GAAGTCATTCCGGTGTGAGC
SEQ ID NO: 27	PKSR4	GCTCTGGATTGGGTATCGCAC
SEQ ID NO: 28	PKSF5	ACTCGAGCATCCGTCGCAATG
SEQ ID NO: 29	PKSR5	AGCAACATCTCCGTCTGGAG
SEQ ID NO: 30	PKSF6	GTTGAACTGTCCATGGCTGA
SEQ ID NO: 31	PKSR6	GAATGAAGGCAATCTGCTGC

All the reactions were carried out in 25- $\mu$ l volumes containing 2.5  $\mu$ l of 10X PCR buffer, 5  $\mu$ l of Q solution, 1  $\mu$ l of each primer (10  $\mu$ M), 0.5  $\mu$ l of 10mM dNTPs mix, and 1.25 U of Taq polymerase (Qiagen Inc., Mississauga, ON, CA). Reaction mixtures were mixed gently and were given flash spin prior to PCR on a Eppendorf Master Cycler<sup>TM</sup> (ep gradient S). PCR amplicons were purified using the QIAquick<sup>®</sup> PCR purification kit (Qiagen Inc., Mississauga, ON, CA; QIAquick is a registered trademark of Qiagen GmbH Corp., Hilden, Fed. Rep. Germany) and commercially sequenced (Plant Biotechnology Institute, Saskatoon, SK).

*Development of a real time RT-PCR assay for detection and quantification of the aur gene in Fusarium species:* All sequences were sequenced, sequences were aligned, several primer sets were designed and tested. One set, named *Auro RT* (SEQ ID NO: 32) and *Auro RTR* (SEQ ID NO: 33) was selected based on efficiency to amplify aurofusarin gene.

SEQ ID NO: 32      ACCTCACTGGAATCAGAGCGCAGC

SEQ ID NO: 33      ATGACRACTTCCCGTGGRCC



The specificity of this set of primers was tested by conventional PCR assay using genomic DNA from *Fusarium* and mycoparasites using the amplification reactions volumes indicated above. The amplification protocol was 1 cycle of 120 s at 94 °C, 35 cycles of 30 s at 94 °C (denaturalization), 30 s at 56 °C (annealing), 45 s at 72 °C (extension), and 1 cycle of 10 min at 72 °C.

The pair of primers used to amplify the  $\beta$ -tubulin (*tub*) gene (the endogenous control gene used to normalize the results) were *FGtubf* (SEQ ID NO: 34) and *Fgtubr* (SEQ ID NO: 35).

SEQ ID NO: 34      GGTCTCGACAGCAATGGTGTT

SEQ ID NO: 35      GCTTGTGTTTTTCGTGGCAGT

The PCR efficiencies of the real time RT-PCR for both genes were checked by performing a 10-fold serial dilution of positive control template to generate a standard curve, and by plotting the Ct as a function of log [10] of template.

Presence of aurofusarin was tested and quantified by RT-PCR in pathogenic *Fusarium graminearum*, *F. graminearum* (3ADON), *F. graminearum* (15ADON), *F. avenaceum*, *F. culmorum*, Mycoparasite, *F. proliferatum*, *F. oxysporum*, *F. arthrosporoides* and mycoparasitic, *Sphaerodes mycoparasitica* and *Trichoderma harzianum*. Relative gene expression was studied when *Fusarium* isolates were co-cultured with Folicur<sup>®</sup>, *Sphaerodes* and *Trichoderma*.

The designed RT-PCR primer set was checked for its specificity by conventional PCR on tested fungal species. A single band 157 bp long was amplified in *F. graminearum* (3ADON), *F. graminearum* (15ADON), *F. avenaceum*, *F. culmorum* but not in other strains. One week old grown cultures tested for their aurofusarin relative gene expression and  $\beta$ -tubulin gene was used as an internal control. *F. graminearum* which produces 15ADON showed highest level gene expression followed by *F. graminearum* (3ADON) and similar kind of expression results were observed with *F. avenaceum* and *F. culmorum*. Aurofusarin relative gene expression was quantified when each *Fusarium* species was co-cultured for a week with *Sphaerodes*, *Trichoderma* and Folicur<sup>®</sup> (Figure 31). Among all tested, *S. mycoparasitica* was most effective in reducing the aurofusarin relative gene expression followed by *Trichoderma* and Folicur<sup>®</sup>

(Figure 31). *S. mycoparasitica* was most affective on *F. graminearum* producing 15ADON, but also efficiently reduced the expression level in *F. culmorum* and *F. avenaceum* and *F.graminearum* producing 3ADON (Figure 31). Similarly, *Trichoderma* was able to somewhat affect *F. graminearum* producing 15ADON and *F. culmorum*, but showed no impact on *F. avenaceum*, whereas enhanced the expression in *F.graminearum* producing 3ADON (Figure 31). Finally, Folicur<sup>®</sup> could be able to provoke minor reduction in *F. graminearum* producing 15ADON *aur* gene expression, but with less impact on *F. culmorum* and *F. avenaceum*. Inversely, Folicur<sup>®</sup> increased expression almost 2-fold when compared to *F. graminearum* producing 3ADON (Figure 31).

*RNA isolation, reverse transcription and real time RT-PCR:* All the cultures were grown for a week under dark and used for expression studies. Fungal total RNA was isolated using the “Total Quick RNA Cells and Tissues” Kit<sup>™</sup> (Bio-Rad Laboratories Inc., Mississauga, ON, CA), according to the manufacturer's instructions, and stored at  $-80^{\circ}\text{C}$ . DNase I treatment to remove the chromosomal DNA contamination from the samples was performed using the “Deoxyribonuclease I, Amplification Grade” (Invitrogen Corp., Carlsbad, CA, USA). First strand cDNA was synthesized using the “Iscript RNA PCR Reagent Kit” (Bio-Rad Laboratories Inc., Mississauga, ON, CA). Relative Quantification of *aur* gene expression was performed in a MiniOpticon Sequence Detection System using the SYBR Green PCR Master Mix (Bio-Rad Laboratories Inc., Mississauga, ON, CA) and the primer pairs indicated above. The PCR thermal cycling conditions for both genes were as follows: an initial step at  $95^{\circ}\text{C}$  for 10 min and 40 cycles at  $95^{\circ}\text{C}$  for 15 s and at  $60^{\circ}\text{C}$  for 1 min. SYBR green PCR master mix 12.5  $\mu\text{l}$  (Bio-Rad Laboratories Inc., Mississauga, ON, CA) was used as the reaction mixture, with the addition of 6.5  $\mu\text{l}$  of sterile milli-Q water, 1.0  $\mu\text{l}$  of each primer (5  $\mu\text{M}$ ), and 5  $\mu\text{l}$  of template cDNA, in a final volume of 25  $\mu\text{l}$ . In all the experiments, appropriate negative controls containing no template were subjected to the same procedure to exclude or detect any possible contamination or carryover. Each sample (triplicate) was amplified twice in every experiment. The results were normalized using the all fungi cDNA amplifications run on the same plate. The *tub2* gene is an endogenous control that was used to normalize quantitation of mRNA target for differences in the amount of total cDNA added to each reaction. Real Time RT-PCR analysis is based on the threshold cycle (CT), which is defined as the first amplification cycle at which the fluorescence

signal is greater than the minimal detection level, indicating that PCR products become detectable. Relative quantitation was the analytical method used in this study. A comparison within a sample is made with the gene of interest (*aur*) to that of the endogenous control gene (*tub2*). Quantitation is relative to the control gene by subtracting the CT of the control gene (*tub2*) from the CT of the gene of interest (*aur*) ( $\Delta$ CT). Each  $\Delta$ CT value (corresponding to each sample) was subtracted by the calibrator value (FpMM6-1C) to obtain the corresponding  $\Delta\Delta$ CT values.  $\Delta\Delta$ CT values were transformed to log<sub>2</sub> (due to the doubling function of PCR) to generate the relative expression levels.

In total 9 isolates were subjected to RT-PCR analyses for quantitative gene expression analyses. Three kinds of colored samples were used in this study composed of 3 white colored, 3 moderately red colored and 3 dark red colored isolates. White colored isolate was used as control and their expression was normalized and taken as one. As reported moderately red colored isolates were able to tolerate 40<sup>0</sup>C whereas; dark red isolates were able to tolerate 80<sup>0</sup>C. When RT-PCR results observed dark colored isolates expressed almost more than 9 folds enhanced relative gene expression whereas, moderately red colored isolates expressed 4 folds enhanced expression (Figure 32).

#### Example 11

Greenhouse studies were performed to assess the effects of *S. mycoparasitica* inoculants on development of *Fusarium* head blight symptoms and the accumulation of tricothecene mycotoxin gene in wheat and barley spikes.

*Fungal isolates and growth:* *S. mycoparasitica* SMCD 2220, and *F. graminearum* 3-ADON chemotype SMCD 2243 were obtained from the Saskatchewan Microbial Collection and Database (SMCD). These fungal cultures were grown on potato dextrose agar (PDA, BD Biosciences, Mississauga, ON, CA) supplemented with antibiotics prior to the study. Mycelial suspensions of these fungal strains (10<sup>4</sup> for *F. graminearum* and 10<sup>6</sup> for both biotrophic mycoparasite and antagonistic fungus) were produced for greenhouse experiment, as follows.

*F. graminearum* 3-ADON chemotype strain was inoculated in potato dextrose broth (PDA, BD Biosciences, Mississauga, ON, CA), incubated at room temperature (~21°C) in

darkness using shaker (100rpm), for a week prior to harvesting the mycelia for greenhouse application. Mycelia of *F. graminearum* were transferred into sterilized commercial blender for cutting the mycelial inoculants into small debris and these mycelia were adjusted to concentration of  $10^4$  CFU/mL.

*S. mycoparasitica* was inoculated in the best preferment yeast peptone dextrose (YPD) (yeast, 5g; peptone, 10g; dextrose, 10g per 1 L of sterile-double distilled water) broth, incubated at room temp in darkness, using shaker at 100 rpm for a week prior to harvesting the mycelia. Mycelia of *S. mycoparasitica* were transferred into sterilized commercial blender for cutting the mycelial inoculants into small debris and the mycelia were adjusted to concentration of  $10^4$  CFU/mL for greenhouse application.

*Greenhouse trials:* The *Fusarium*-susceptible wheat CDC-TEAL and barley BOLD cultivars were used to test the efficacy of biotrophic mycoparasite *S. mycoparasitica* and one potential *Fusarium* antagonistic fungal isolate as potential biocontrol agent for managing *Fusarium* head blight diseases. Surface-sterilized wheat and barley seeds were planted in 15-cm diameter pots containing Pro-Mix<sup>®</sup> soil (Sun Gro Horticulture, Delta, BC, CA) and maintained at 23-25 °C during the day time and 18-20 °C during the night time in a greenhouse with light intensity of  $360 \mu\text{mol m}^{-2} \text{s}^{-1}$ , watered every 2 d, and fertilized every 14 d using 1300 ppm of NPK (20-20-20) fertilizer. All treatments in the experiment were with three replicates and repeated twice.

At the anthesis stage, wheat and barley spikes were sprayed with mycelial suspension of *S. mycoparasitica* at a concentration of  $10^6$  CFU/mL, antagonistic fungal isolate at a concentration of  $10^6$  CFU/mL, 65  $\mu\text{mol/L}$  tebuconazole, or sterile distilled water. The treatments were applied as outlined in Xue et al. (2009, Can. J. Plant Pathol. 31: 169-179) with slight modifications. Following the inoculation of potential biocontrol agents, plants were kept at room temperature and covered with sterile Whirl-Pak<sup>®</sup> bags (Whirl-Pak is a registered trademark of Aristotle Corp., Stamford, CT, USA) for overnight to allow fungal growth prior to *F. graminearum* inoculation. Plants were then inoculated with mycelial suspension of *F. graminearum* at a concentration of  $10^4$  CFU/mL. *Fusarium graminearum* inoculum was sprayed on the spikes and covered with sterile Whirl-Pak<sup>®</sup> bags for overnight. The inoculated plants were maintained for an additional 21 days prior to sampling. Percentage of infected spikelets (IS) per

spike, FHB index, FDK, and weight of 100 seeds were rated as outlined in Xue, A.G., Voldeng, H.D., Savard, M.E., Fedak, G., Tian, X., and Hsiang T. (2009) Biological control of fusarium head blight of wheat with *Clonostachys rosea* strain ACM941, Canadian Journal of Plant Pathology 31: 169–179. The FHB visual severity scale was determined.

The biocontrol effects of different concentrations of *S. mycoparasitica* of *Fusarium* head blight symptoms in barley are shown in Figure 33. The data in Figures 33A, 33B, and 33C show that inoculation of a susceptible barley cultivar with *F. graminearum* significantly reduced that height of the plants, average numbers of spikes form per plant, and the average weight of 5 spikes. However, inoculation of the barley cultivar with *S. mycoparasitica* did not affect growth and development. Treatment of *F. graminearum*-infected barley with *S. mycoparasitica* at an inoculation level of  $10^6$  CFU/mL prevented the onset of any *Fusarium* head blight symptoms. Treatment of *F. graminearum*-infected barley with lower concentrations of *S. mycoparasitica*, i.e.,  $10^4$  CFU/mL and  $10^5$  CFU/mL, did not protect against the occurrence of the disease symptoms (Figures 33A - 33C).

The biocontrol effects of different concentrations of *S. mycoparasitica* of *Fusarium* head blight symptoms in wheat are shown in Figure 34. The data in Figures 34A, 34B, and 34C show that inoculation of a susceptible wheat cultivar with *F. graminearum* significantly reduced that height of the plants, average numbers of spikes form per plant, and the average weight of 5 spikes. However, inoculation of the wheat cultivar with *S. mycoparasitica* did not affect growth and development. Treatment of *F. graminearum*-infected wheat with *S. mycoparasitica* at an inoculation level of  $10^6$  CFU/mL prevented the onset of any *Fusarium* head blight symptoms. Treatment of *F. graminearum*-infected wheat with lower concentrations of *S. mycoparasitica*, i.e.,  $10^4$  CFU/mL and  $10^5$  CFU/mL, did not protect against the occurrence of the disease symptoms (Figures 34A - 34C).

The data in Figure 35 and Table 8 show that treatment of *F. graminearum*-infected barley with *S. mycoparasitica* at an inoculation level of  $10^6$  CFU/mL provided a comparable level of protection against the occurrence of *Fusarium* head blight symptoms, as was provided by the commercial fungicide Folicur<sup>®</sup>.

Table 8: Effect of *S. mycoparasitica*, Follicur® fungicide and *F. graminearum* on percentage of infected spikelets, *Fusarium* head blight index, percentage of *Fusarium* damaged kernels, and weight of 100 seeds of the barley cultivar BOLD in greenhouse trials.

Treatment	Infected spikelets (%)	<i>Fusarium</i> head blight index (%)	<i>Fusarium</i> damaged kernels (%)	Weight of 100 seeds (g)
<i>F. graminearum</i> (Fg)	62.6±8.0 a	56.7±11.9 a	75±9.4 a	4.1±0.4 c
<i>S. mycoparasitica</i> – Fg	15.4±3.1 bc	4.1±1.0 b	19±6.5 b	5.4±0.6 ab
Follicur® – <i>F. g</i>	19.6±6.9 b	7.2±4.8 b	16±6.5 b	5.7±0.5 a

*Real-time PCR quantification:* Total DNA was extracted from the wheat and barley spikes with DNeasy<sup>®</sup> Plant Mini Kit (Qiagen Inc., Mississauga, ON, CA). Extracted total DNAs from spikes of different treatments were employed in real-time PCR quantification. Two different primer sets and PCR conditions used in this study were described in Nicholson et al. (1998) for Fg16NF/R primer set and Schnerr et al. (2001) for Tox5-1/2 primer set. Total DNA extracted from spring wheat and barley spikes harvested from greenhouse trials were carried out in a MiniOpticon (Bio-Rad Laboratories Inc., Mississauga, ON, CA). All real-time PCR reactions were performed using real-time PCR MJ white tubes (Bio-Rad Laboratories Inc., Mississauga, ON, CA) with a total volume of 25 µl of IQ supermix (Bio-Rad Laboratories Inc., Mississauga, ON, CA), 1 µl of each 10 µM forward/reverse primers (Invitrogen), 3.4 µl of BSA (Bovine Serum Albumin) (1.47 µg/µl) (Ishii, S. and Loynachan, T.E. (2004) Rapid and reliable DNA extraction techniques from trypan-blue-stained mycorrhizal roots: comparison of two methods. *Mycorrhiza* 14:271-275), 5.1 µl of sterilized UltraPure Millipore water, and 2 µl of DNA template.

Figure 36 shows standard curves of *F. graminearum* chemotype 3 genomic DNA concentration standards versus cycle threshold (Ct) with PCR reactions performed in triplicate using primer sets (A) *Tox5-1/2*, with genomic DNA ranging from 270 ng ( $\text{Log}_{10} = 2.90$ ) to 0.27 ng ( $\text{Log}_{10} = -0.60$ ), readings at 0.005 fluorescence line and (B) *Fg16NF/R*, with DNA template ranging from 270 ng ( $\text{Log}_{10} = 2.43$ ) to 0.027 ng ( $\text{Log}_{10} = -1.57$ ); in 10-fold dilution series, readings at 0.025 fluorescence line. Error bars indicate standard deviation for the mean of *F. graminearum* chemotype 3 standard curves derived from *tri5* gene and *F. graminearum* specific primer sets. Figure 37 shows effects of *S. mycoparasitica* (B) and Folicur fungicide (Fol) treatments on *F. graminearum* chemotype 3-ADON genomic DNA detected in barley spikes employing RT-PCR. Treatments were: Fus – *F. graminearum*; B-Fus – *S. mycoparasitica* with *F. graminearum*; Fol-Fus – Folicur fungicide with *F. graminearum*. All values obtained were the means of six replicates. Error bars indicate standard deviation of the mean. Means of *F. graminearum* DNA with Tox 5 primer were log<sub>10</sub> transformed prior to LSD test. Both primer sets were analyzed separately. Values followed by the same letters within each primer set are not significantly different using LSD test at  $P < 0.05$ .

## Example 12

In addition to the known *Fusarium* spp. hosts for *S. mycoparasitica* (i.e., *F. avenaceum*, *F. graminearum* and *F. oxysporum*), we surprisingly discovered that *F. culmorum* and *F. equiseti* are also mycoparasitised by *S. mycoparasitica* as disclosed in the following study.

*Fungal isolates and growth:* Biotrophic mycoparasites *S. mycoparasitica* SMCD2220, *Sphaerodes quadrangularis* strain CBS112764, *Sphaerodes retispora* var. *retispora* strain CBS 994.72, and pathogenic *Fusarium* strains (*F. arthrosporioides* SMCD2247, *F. culmorum* SMCD2248, *F. equiseti* SMCD2134, *F. flocciferum* SMCD2135, *F. poae* SMCD2136, and *F. torulosum* SMCD2139) were obtained from the Saskatchewan Microbial Collection and Database (SMCD), Saskatchewan, Canada. All fungal isolates were grown and maintained on potato dextrose agar (PDA, BD Biosciences, Mississauga, ON, CA) prior to the study.

*Fungal-fungal interactions:* For examination of the interaction between isolates of *Sphaerodes* and *Fusarium* species, both biotrophic mycoparasite and *Fusarium* isolates were inoculated and assessed using slide culture assays. Slides were maintained in a sterile humidity chamber and daily observations on the hyphal interactions at the meeting place (contact zone) were performed under a Carl Zeiss Axioskop2 equipped with Carl Zeiss AxioCam ICc1 camera with 20×, 40× and 100× objectives. Formation of biotrophic mycoparasitic contact structures attaching *Sphaerodes* species to *Fusarium* hyphae were examined, recorded and compared to drawings from the literature (Jordan & Barnett, 1978, *Mycologia* 70: 300-312; Rakvidhyasastra & Butler, 1973, *Mycologia* 65: 580-593; Whaley & Barnett, 1963, *Mycologia* 55: 199–210). Diameters of both parasitized and non-parasitized *Fusarium* hyphal cells were measured under light microscopy with a 100× objective lens. Each treatment used six replications consisted of *Sphaerodes* or *Fusarium* alone, and *Sphaerodes*–*Fusarium* co-inoculated. The experiment was repeated twice. In the slide-culture assay, *Fusarium* mycelia infected with *Sphaerodes* haustoria were stained with lactofuchsin. Stained hyphae of both *Fusarium* and *Sphaerodes* in slide-culture were then examined with a Carl Zeiss Axioskop2 fluorescent microscope attached to Carl Zeiss AxioCam ICc1 with 40× and 100× objectives. Slide-culture assays were also subjected to Zeiss META 510 confocal laser scanning microscopy (CLSM) analysis to observe intracellular mycoparasitism under a C-Apochromat 63× N.A.1.2 phase-contrast water immersion objective



through Z-stacking mode to scan through the *Fusarium* hyphae with intracellular infection (CLSM with 514nm excitation – argon and LP585 emission filters).

Hyphae-hyphae interactions and contact structures in the contact zone were examined for seven days. On day three, *Sphaerodes mycoparasitica* was found to produce hook-shaped contact structures on *F. equiseti* and *F. culmorum* (Figure 39). On day five, more hook-shaped contact structures and intracellular penetration of *F. equiseti* were observed (Figures 39A, 39A-39D). The combination of lactofuchsin dye and fluorescent or confocal laser scanning microscopy revealed that the parasitized or penetrated *Fusarium* cells became empty (loss of cytoplasm = no fluorescence) or fluoresced with low intensity (very pale) (Figures 39A-39D) as compared to healthy *Fusarium* cells. During the seven days of observation, no *S. mycoparasitica* hyphae were observed within *F. culmorum* cells. *S. mycoparasitica* produced hook-shaped contact structures (Figure 38A, a) more frequently than clamp-like contact structures (Figure 38B, b) on both *F. equiseti* and *F. culmorum*. Diameters of *F. equiseti*, but not *F. culmorum*, hyphae parasitized by *S. mycoparasitica* were observed to be significantly reduced compared to non-parasitized *Fusarium* hyphae (with T-test,  $P = 0.001$  and  $P > 0.05$ , respectively) (Figure 41).

None of the *Fusarium* taxa tested appeared to be suitable hosts for mycoparasitic *S. quadrangularis* and *S. retispora* even after 10 days of co-inoculation on slide cultures. No contact biotrophic parasitic structures or intracellular parasitism by *S. quadrangularis* and *S. retispora* on the tested *Fusarium* strains were observed at the interaction or contact zone. Also, *F. arthrosporioides*, *F. flocciferum*, *F. poae*, and *F. torulosum* did not appear to be suitable hosts for *S. mycoparasitica*. Around five days after inoculation on slide culture assays, mycelia of *F. arthrosporioides* were inhibited by *S. mycoparasitica*. *F. arthrosporioides* started to form rosette-like mycelia at the contact zone with *S. mycoparasitica* (Figure 39B).

On the fifth and seventh days after inoculation, anamorphic structures were produced by *S. mycoparasitica* more abundantly in the zone of contact with *F. culmorum* (Figures 40C and 40D). Anamorphic structures or asexual organs in close proximity to *F. culmorum* mycelia were red-colored (Figure 40D), whereas the organs at a distance were not (Figure 40C).

**CLAIMS**

1. An isolated culture of *Sphaerodes mycoparasitica*, wherein the species is characterized by a combination of:

(a) fusiform and triangular ascospore shape, and reticulate and smooth wall ornamentation;

(b) conidia produced from simple phialides on the surface of ascoma peridial wall on ascoma surrounding hyphae, and on irregularly branched conidiospores arising from hyphae; and

(c) forming hook-like structures parasitizing living hyphae of *Fusarium*;

wherein the isolated culture of *Sphaerodes mycoparasitica* comprises a gene encoding a large subunit of ribosomal RNA gene as shown in SEQ ID NO:1; a gene encoding a small subunit of ribosomal RNA as shown in SEQ ID NO:2; and a gene encoding an internal transcribed spacer ribosomal DNA as shown in SEQ ID NO:3.

2. The isolated culture of *Sphaerodes mycoparasitica* according to claim 1, wherein the isolated culture is *Sphaerodes mycoparasitica* strain deposited as IDAC 301008-01.

3. A use of the isolated culture of *Sphaerodes mycoparasitica* of claim 1 or 2 for controlling *Fusarium* spp.

4. The use of claim 3, for controlling *Fusarium* spp. in plants.

5. A use of the isolated culture of claim 1 or 2, for modulating synthesis of or degrading one of a *Fusarium* trichothecene mycotoxin deoxynivalenol (DON), mycotoxin 3-acetyldeoxynivalenol, mycotoxin 15-acetyldeoxynivalenol, mycotoxin zerelanone, and mycotoxin aurofusarin.

6. A composition comprising the isolated culture of *Sphaerodes mycoparasitica* according to claim 1 or 2 and a carrier.

7. The composition of claim 6, further comprising an additional antifungal agent.

8. A method for controlling *Fusarium* spp. in plants, the method comprising treating a batch of seeds with the culture of claim 1 or 2 or the composition of claim 6 or 7 and then culturing the treated seeds into plants.
9. A method for controlling *Fusarium* spp. in plants, the method comprising treating growth medium contaminated with the *Fusarium* spp. with the culture of claim 1 or 2 or the composition of claim 6 or 7.
10. The method of claim 9, wherein the growth medium is soil.




Figure 1




Figure 2




Figure 3






Figure 4



(a)



(b)

Figure 5






Figure 6





Figure 7



A

B

Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13






Figure 14




Figure 15




Figure 16




Figure 17




Figure 18

19/41





Figure 19




Figure 20




Figure 21






Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29






Figure 30




Figure 31






Figure 32

A: Height of plants (cm)



B: Spikes per plant (average number)



C: Weight of 5 spikes (g)






Figure 33

A: Height of plants (cm)



B: Spikes per plant (average number)



C: Weight of 5 spikes (g)




Figure 34




Figure 35




Figure 36






Figure 37





b

a

Figure 38



Figure 39




Figure 40




Figure 41