



Sustainable Environment

An international journal of environmental health and sustainability

ISSN: (Print) (Online) Journal homepage: www.tandfonline.com/journals/oaes21

Climate smart agriculture strategies for enhanced agricultural resilience and food security under a changing climate in Ethiopia

Abdulbasit Hussein |

To cite this article: Abdulbasit Hussein | (2024) Climate smart agriculture strategies for enhanced agricultural resilience and food security under a changing climate in Ethiopia, Sustainable Environment, 10:1, 2345433, DOI: [10.1080/27658511.2024.2345433](https://doi.org/10.1080/27658511.2024.2345433)

To link to this article: <https://doi.org/10.1080/27658511.2024.2345433>



© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 26 Apr 2024.



Submit your article to this journal [↗](#)



Article views: 229



View related articles [↗](#)

Climate smart agriculture strategies for enhanced agricultural resilience and food security under a changing climate in Ethiopia

Abdulbasit Hussein 

Department of Natural Resource Management,, Haramaya University College of Agriculture and Environmental Science (HU CAES), Dire Dawa, Ethiopia

ABSTRACT

Climate Smart Agriculture (CSA) is a practice aimed at increasing agricultural productivity and income while reducing greenhouse gas emissions and strengthening agricultural resilience against climate change. In Ethiopia, the agricultural industry is being impacted by climate change, and various strategies are being implemented to enhance sustainability and food security. These include promoting climate-resilient crop breeds, integrating conservation agriculture techniques, using advanced livestock management practices, and establishing effective water management systems. CSA programs also aim to improve access to credit, extension services, and markets for smallholder farmers. However, challenges such as limited finance, technical capacity, and inadequate infrastructure hinder the adoption and scaling up of these strategies. Despite these challenges, CSA is a crucial step towards achieving food security and can serve as a model for other countries facing similar challenges.

ARTICLE HISTORY

Received 28 September 2023
Accepted 16 April 2024

KEYWORDS

Ethiopia; climate change; strategies; adaptation; mitigation

1. Introduction

1.1. Backgrounds of climate change and agriculture in SSA

Climate change is attributed to the escalation of greenhouse gas emissions, namely nitrous oxide (N₂O), carbon dioxide (CO₂), and methane (CH₄), resulting in erratic precipitation cycles, elevated temperatures, droughts, and floods (Jogdand, 2020; H. Zegeye, 2018). African countries are among the most susceptible areas globally, and the results of this susceptibility are anticipated to be severe, widespread, and enduring in the coming years. Numerous recent studies have revealed the vulnerability of African countries to climate change. According to Hulme et al. (2001) and the IPCC (2014), by 2050, the region of East Africa will experience elevated temperatures, along with a notable increase in rainfall between December and February, ranging from 5–20%. On the other hand, a 5–10% decrease in rainfall is anticipated from June through August. Climate change will lead to a decline in food production, particularly in maize production, as well as adverse effects on aquaculture and coastal fisheries (Adhikari et al., 2015). Furthermore, drought and an increase in the mean annual temperature were shown

to be the most common climatic conditions in sub-Saharan Africa, which pose a considerable danger to crop production systems dependent on rainwater and the livelihoods of subsistence farmers (Zougmore et al., 2016).

According to Allen et al. (2020), the agriculture sector is responsible for between 10 to 20% of all anthropogenic greenhouse gas emissions. Because climate change negatively affects crop, livestock, and fishery production yields, it poses a danger to food and nutrition supply. Population growth (Molyneux et al., 2012), urbanisation (Srivastava & Srivastava, 2020), mechanisation (Kolberg et al., 2019), and globalisation (McMichael, 2013) will also exacerbate and hasten these effects. Adoption of sustainable systems in food systems, general consumption patterns, and agriculture has been suggested in a number of technical studies and study findings (Powlson et al., 2014). By reducing greenhouse gas emissions and the carbon footprint, switching to renewable energy sources from non-renewable ones, conserving genetic resources, maintaining regional breeds that are adapted to their particular environments, and fostering biodiversity, this recommendation seeks to lessen the negative effects of these activities (Khelifa et al., 2021; Mohamed-Brahmi et al., 2022).

CONTACT Abdulbasit Hussein  abdulbasithussein111@gmail.com  Department of Natural Resource Management, Haramaya University College of Agriculture and Environmental Science (HU CAES), P. O Box 138, Dire Dawa, Ethiopia

Reviewing editor: Tessema Toru Department of Natural Resource Management, Haramaya University College of Agriculture and Environmental Science (HU CAES), Dire Dawa, Ethiopia

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

According to FAO (2013), CSA refers to farming methods that support resilience while also encouraging sustainable productivity, reducing greenhouse gas emissions when practical, and helping countries meet their goals for food security and development (Campbell, 2017; Lipper et al., 2014). It has been acknowledged that the application of CSA is an essential tool for tackling the challenges that climate change presents to agricultural systems and for more successfully integrating agriculture into international climate discussions (Andrieu et al., 2021). As a matter of fact, CSA assists major institutions, service providers, and farmers in building their capacity to handle the risks associated with rising climatic variability as well as adapt to long-term climate change (Zougmore et al., 2016). The agricultural industry can help reduce the effects of climate change and boost resilience by implementing CSA (Williams et al., 2015). Additionally, CSA strategies seek to mitigate and adapt to the effects of climate change, as well as provide smallholder farmers and vulnerable people with a fair and consistent income and favorable working situations (Westermann et al., 2018). Africa's agricultural diversity, fishing practices, and technologies are complemented by CSA methods and technologies that offer integrated possibilities. These integrated solutions are essential for adapting to climate change and include agroecological methods, ecosystem management, and sustainable natural resource management (Nyasimi et al., 2014).

CSA methods can significantly improve local economies and social justice, especially in African nations (Mutenje et al., 2019; Taylor, 2018; Westermann et al., 2018). The increased interest in community support agriculture (CSA) has led to a number of parties launching various CSA initiatives in recent years, including farmers, governments, international organisations, civil society organisations, the scientific community, and the commercial sector (Dinesh et al., 2015). In Ethiopia, climate-smart agriculture was first introduced and put into practice more than 10 years ago by the government and many NGOs, including SOS Sahel, Farm Africa, Climate Change Forum, Self Help Africa, SG2000, CARE, and World Vision. Building farmers' competence and giving them the authority to implement CSA measures has been stressed as a means of sustaining agricultural activities in SSA, especially in Ethiopia (Djenontin et al., 2004). Because agricultural practices are influenced by demographic, institutional, economic, and physical factors, the results of previous studies on

climate-smart agriculture have been inconsistent (Belay et al., 2023; Ebabu & Okoyo, 2017; Fentie & Beyene, 2019; Tekeste, 2021; Wassie & Pauline, 2018; Wekesa et al., 2018). This review aims to develop climate-smart agriculture strategies in Ethiopia to enhance community livelihoods and promote biodiversity conservation.

2. Methodology

2.1. Literature search method

The current review study included a variety of search subjects to locate scholarly articles related to climate smart agriculture strategies and food security research that benefits local communities in Ethiopia. To the greatest extent possible, the scope and research areas within the subject matter were covered by the broad definition of the search themes. The pertinent scientific publications were gathered from the core collection databases of the Web of Science (WoS), Scopus, and Google Scholar. Documents were searched by inputting a combination of keywords: (CSA), (strategies) and (impacts of climate on food security), as well as with 'Ethiopia', for areal restriction. Although the study region was confined to Ethiopia. The documents that were retrieved included core details including authors, titles, keywords, abstract texts, nations, institutions, publications, and referenced references. The selection adhered to the Preferred Reporting Items for Systematic Reviews as the standard for the collection, choice, analysis, and reporting of journal articles. The literature selection procedure was provided as a PRISMA flow diagram in 2020).

A preliminary examination was conducted by perusing the titles and abstracts of the papers to confirm their eligibility for inclusion. Based on a scrutiny of published synopses, reviews, letters to the editor, book chapters, conference abstracts, and research articles that (1) were published before 2010, (2) recorded data from countries other than Ethiopia, (3) recorded data from intensive agriculture, or (4) covered scientific and technical reports that were not published in English were excluded. Further, only comprehensive articles were retained, and duplicative articles were expunged. After an exhaustive study of the chosen publications, the requisite data encompassing the first author, research year, provincial and geographic information were extracted. Overall, 152 documents were considered for this review paper.

3. The application of CSA practices and technologies that have been adopted and implemented in Ethiopia

A wide range of agricultural development programmes have been implemented in Ethiopia, utilising both traditional and innovative methods to guarantee food supply and enhance sustenance (Ntawuruhunga et al., 2023). Currently, the country's agricultural development operations are supported by a number of policies, initiatives, and institutions (Berhanu & Poulton, 2014). The numerous agricultural development programmes that have been implemented should be evaluated to determine which are most important for tackling climate change-related issues and supporting their adaptation and mitigation (Conway & Schipper, 2011). Some of the agricultural practices used in Ethiopia include integrated watershed management, composting, rangeland management, conservation agriculture, sustainable land management, crop residue management, agroforestry, and the promotion of better animal feed (Figure 1).

Ethiopia has seen a rise in the use of Climate-Smart Agriculture (CSA) technologies and practices as part of initiatives to mitigate the effects of climate change and advance sustainable agricultural systems. The following are some instances of CSA methods and tools that Ethiopia has embraced and put into use:

3.1. Integrated watershed management

In Ethiopia, integrated watershed management is conducted through various projects and programs, namely sustainable land management programs (SLMP1 and SLMP2) (Adimassu et al., 2016), the Managing Environmental Resources to Enable Transitions to More Sustainable Livelihoods (MERET) project (Gashaw, 2015), Productive Safety Nets Programme Public Works (PSNP-PW) (Bahru & Zeller, 2022), and several non-governmental organizations. The SLMP2 program employs advanced techniques like 'climate-smart agriculture', which involves the use of drought-resistant food crops, alongside traditional methods such as mulching, intercropping, conservation agriculture, no-till, crop rotation, cover cropping, integrated crop-livestock management, agroforestry, improved grazing, and integrated crop-livestock management (Zerssa et al., 2021). Reports indicate that the program has resulted in the closure of approximately 1,708,100 ha of land and the application of appropriate biological and physical soil conservation methods on an additional 2,076,000 ha of land in various regions of the country.

While maize is the main crop grown in the Debremawi Watershed, teff, barley, faba bean, and haricot bean are all important crops grown there (Tilahun et al., 2014). The low agricultural output in the watershed is caused by a variety of factors, such as severe soil erosion, decreased soil fertility, complete removal of




Figure 1. A diagram of the various routes to CSA. Source: (Were et al., 2016)

crop residue for use as fuelwood and livestock feed, and inadequate extension services (Zegeye et al., 2016). To address this issue, the Amhara Region Natural Resource Management Bureau and the Water and Land Resource Centre (WLRC) collaborated to initiate an intervention in 2012. According to Tesfaye et al. (2018), this non-governmental organisation (NGO) has made great progress in resolving the issue at hand. Preventing environmental degradation, increasing agricultural output, and enhancing population food security were the objectives. To this end, the watershed is implementing improved animal management practices, improved crop types, and physical soil protection (Tilahun et al., 2014).

The ecosystem of the area has undergone rapid transformation as a result of the enclosure of degraded areas and the avoidance of open grazing. Degraded plains were covered with trees and grass within two years, and gullies began to replenish. Open grazing is discouraged; therefore, some crop residues are left on the soil, increasing the amount of organic matter. Soil erosion was greatly reduced, and water infiltration was greatly improved. Crop types compatible with local demands and environments have been adopted by farmers. The work done thus far has made it possible for farmers to embrace other CSA practices such as planting fruit trees, small-scale irrigation, and the construction of feedlots (Mhired et al., 2020).

In the Oromia region, 49% of the sample households use these structures, compared to 44% in the SNNP region, according to the findings on the use of various soil and water conservation structures (Diro et al., 2022). Therefore, conservation structures should be used to prevent termites and reduce soil acidity. Therefore, farmers in Oromia adopt more stringent conservation measures than those in the SNNP region.

Diro et al. (2022) further, reported the households that have adopted physical soil and water conservation measures, 45% employ soil bunds, with 53 percent doing so in Oromia and 37% doing so in the SNNP region. Farmers in the study areas also used terracing and stone bunds after using soil bunds (Figure 2).

3.2. Integrated soil fertility management

In order to improve agricultural output and solve the urgent issue of food insecurity, the Ministry of Agriculture has focused its attention on controlling soil fertility (Hörner & Wollni, 2021). A national initiative, mostly through national and regional extensions, aims to enhance farmers' capacity to control soil fertility (Agegnehu & Amede, 2017). Composting, crop rotation, intercropping, training, and the promotion of appropriate fertiliser application are important programmes targeted at improving food security (Yebo, 2015). As a result, a large number of farmers have adopted intercropping, better fertiliser use, and improved compost preparation and application. Recent evaluations indicate that improved extension services have led to a notable rise in crop productivity in these nations (Hörner & Wollni, 2021).

3.3. Conservation agriculture

Traditionally, farmers in Ethiopia employ minimal tillage and other soil conservation techniques. Nevertheless, the earnest promotion of conservation agriculture technology began in 1998, when Sasakawa Global (SG2000), Makobu, and regional agricultural development bureaus jointly promoted and demonstrated the technology on 77 farmers' plots (FAO,




Figure 2. Type physical soil and water conservation structures used in %. Source: (Diro et al., 2022)

2016). Trials on maize, sorghum, and teff were conducted between 1999 and 2003 at the Jima, Bako, and Melkasa research centers during the early introduction of conservation agriculture. The studies showed that compared to conventional tillage, conservation tillage plots produced higher yields (Burayouet et al., 2002). The studies also showed that the conservation of agricultural lands had reduced production costs. Based on the comprehensive pattern revealed by these data, it can be posited that conservation agriculture exhibits enhanced yields in both the immediate and prolonged terms. This observation is consistent with the analyses of research conducted in Africa, Latin America, and Asia, which revealed that yields from conservation agriculture surpass those from conventional agriculture by a range—20–120 percent (Derpsch et al., 2010; Kassam et al., 2009). Mulching and residue control can improve soil fertility and nutrient availability (Ngosong et al., 2019). Another important strategy for yield stabilization or improvement is to increase water availability throughout the cropping cycle (Altieri et al., 2015).

In cooperation with Federal and Regional Agricultural Offices, the Food and Agriculture Organisation (FAO) offered technical and financial support for the promotion of conservation agriculture in Ethiopia in 2010 (Brown et al., 2017). 24 conservation agriculture demonstration plots were established with the active participation of 600 smallholder farmers from 12 woredas districts in the Amhara, Oromia, and Tigray regions (FAO 2016). In addition, the FAO helped 72 extension agents receive training so they could lead farmer field schools focused on conservation agriculture. Remarkably, 32 of these agents received instruction on the setup and maintenance of equipment used in conservation agriculture. In 2010, the previously stated woredas received a variety of conservation agriculture equipment, such as fertiliser planters, jab planters, and oxen-drawn seeds (FAO, 2016).

According to multiple sources, conservation agriculture has been adopted on a wide scale in regions where it has been firmly established, such as in select areas of Oromia, Amhara, and Tigray (Araya et al., 2011). These specific regions comprise the West Gojam, East Gojam, and South Gonder Zones of the Amhara Regional State, as well as the West, East, South, and West Wollega Zones of the Oromia Regional State. These areas are renowned for the production of teff and maize (FAO 2016).

According to research conducted in the Oromia Region's Bako and Adaa Woreda, more than half of the sample respondents (57.4%) were found to utilize one or more components of the conservation tillage technology package (Liben et al., 2018; Tsegayea et al., 2010). Of the early adopters, 10% have utilized only one

component, 75% have used two components, and 15% have implemented all three components (Liben et al., 2018; Tsegayea et al., 2010). Furthermore, Tsegaye et al. (2015) reported that adopters of three conservation agriculture practices—mulching, limited plowing, and crop rotation—have higher yields than non-adopters. These three elements significantly enhance labor productivity (yield per unit of labor). Farmers in regions where teff is grown have more incentives to use conservation agricultural techniques because teff requires more labor than maize (Tsegaye et al., 2015). The report also stated that crop yields increase with the adoption of more conservation agriculture components and that labor per unit of crop yield decreases as more conservation agriculture components are used. This indicates that the full implementation of all conservation agriculture principles results in labor savings (Araya et al., 2011; Liben et al., 2018; Tsegaye et al., 2015).

Most farmers readily incorporate herbicides into their farming practices in regions where weed issues are common (Mengistie et al., 2017). Teff requires frequent plowing; thus, farmers in Adaa District, where it is necessary to grow the crop and use minimum plowing, are significantly more economical (Tsegayea et al., 2010). Teff requires heavy plowing four to six times before planting; therefore, conservation agriculture considerably reduces labor costs (Araya et al., 2015).

One of the woredas in Oromia Regional State's East Wollega Zone where smallholder farmers use conservation agriculture is Sibu Sire (Merga, 2020). To raise total crop yield in the Woreda, SG2000 Ethiopia started a programme in 2001 (Fayera, 2022). One of the techniques is conservation agriculture. Farmers, Woreda subject matter experts (SMS), and development agents (DAs) get extensive training in conservation agriculture practices such crop rotation, crop residue management, and reduced tillage (FAO, 2016). Farmers received instruction on appropriate weed management practices, with a focus on the proper administration of non-discriminatory herbicides, such as roundup. Numerous displays of maize and teff preservation farming were held, and productive field days for exhibitions were planned (FAO 2016). A report from the Woreda office states that on about 3500 hectares of land, mostly in maize and teff, about 4000 smallholder farmers practiced conservation agriculture in 2014 (FAO 2016). v

3.4. Agroforestry practice in Ethiopia

Agroforestry is a long-standing agricultural practice prevalent in Ethiopia (Amare et al., 2019; Bekele, 2018; Haile et al., 2019; Legesse & Negash, 2021; Lelamo, 2021). This process involves incorporating trees and

bushes into farms through natural regeneration or planting. The cultivation of the moringa tree has a long-standing history in various areas of Ethiopia's Konso, Omo, Burji, Sena, and Mele woredas in the SNNP Regional State, where it is typically grown alongside sorghum and other commodities, as noted by Shonde (2017). The significance of this tree as an essential technology for climate change adaptation has been recognized by the Agricultural Extension Directorate of the MoA, and promotion efforts are currently underway in South and North Wollo, East and West Hararghe, Sidama Zones, and other areas through the establishment of nurseries and seed collection, according to Birhane (2014) and Diriba et al. (2021). Additionally, the Relief Society of Tigray (REST) has advocated agroforestry as an integral component of sustainable agriculture in multiple regions of northern Ethiopia, as stated by Gebru et al. (2019). The Agricultural Extension Directorate has also formulated a technology package for the extensive promotion of *Faidherbia albida* in Ethiopia. As part of the Climate Resilient Green Economy (CRGE) Strategy, the Ethiopian government initiated a nationwide program in 2011 to plant more than 100 million *Faidherbia* trees in farmers' fields (Paul & Weinthal, 2019).

Research has shown that adding shrubs or perennial trees to Ethiopian agricultural fields used for crop production and grazing enhances soil cover and guarantees green cover all year round (Ketema & Yimer, 2014). The Forestry Research Centre (FRC) conducted agroforestry tree on farmon-farm trials in the Meiso and Boset districts of the Oromia Region. The results varied in terms of maize yield when compared to conservation agriculture and conventional tillage (FAO, 2016).

Farmers in Ethiopia have maintained multipurpose trees, such as *Croton macrostachyus*, *Erythrina brucei*, *Albizia gummifera*, *Millettia ferruginea*, and *Cordia africana*, which are important components of agricultural landscapes (Yadessa et al., 2009). *Ekebergia capensis*, *Cordia africana*, *Olea capensis*, *Millettia ferruginea*, *Erythrina brucei*, *Annona senegalensis*, and *Citrus medica* are other significant plants in the country's southern region, particularly in home gardens, where they are handled using farmers' traditional knowledge (Adane et al., 2019; Anshiso et al., 2017; Gina et al., 2014; Mesele, 2007)(Table 1).

3.4.1. The roles of agroforestry trees for livelihood in Ethiopia

To maximise the acquisition and use of resources for the environment, a household often preserves native multifunctional trees on farmland for a number of critical and beneficial functions (Mesele, 2007). This is contingent upon the useful advantages they offer the household, including sustenance, fuel, windbreaks, soil fertility, and a host of other commodities (Abebe et al., 2010; Darcha et al., 2015; Lelamo, 2021; Linger, 2014; Negash & Starr, 2015). The preservation of arboreal taxa is supported by additional justifications that are related to apiculture and other revenue-generating activities (Edo et al., 2017; Muleta et al., 2011). Fruit trees with several uses are primarily used for food, especially in hard droughts. The fruit tree species that are widely used in northwest Ethiopia are indicative of the high level of farmers' reliance on locally grown food in the western Hararge zone of the Oromia region of Ethiopia (Linger, 2014). Furthermore, these trees are used not just for personal use but also to provide revenue through the sale of tree

Table 1. Ethiopian multipurpose agroforestry tree species

Major trees species	Area in Ethiopia	Sources
<i>Citrus medica</i> , <i>Annona senegalensis</i> , <i>Cordia africana</i> , <i>Erythrina brucei</i> , <i>Ekebergia capensis</i> , <i>Millettia ferruginea</i> , <i>Ficus vasta</i> , <i>Prunus africana</i> , <i>Syzygium guineense</i> , <i>Moringa stenopetala</i> , <i>Olea capensis</i> and <i>Vernonia schimperi</i>	Southern part of Ethiopia	(Adane et al., 2019; Agize et al., 2013; Asfaw & Ågren, 2007; Gebretsadik & Negash, 2016; Gina et al., 2014; Negash, 2007)
<i>Cordia africana</i> , <i>Erythrina abyssinica</i> , <i>Croton macrostachyus</i> , and <i>Vernonia amygdalina</i>	Eastern parts of Ethiopia	(Gindaba et al., 2005; Mamo & Asfaw, 2017)
<i>Acacia abyssinica</i> , <i>Albizia schimperiana</i> , <i>Albizia gummifera</i> , <i>Cordia africana</i> , <i>Erythrina abyssinica</i> , <i>Croton macrostachyus</i> , <i>Ficus thonningii</i> , <i>Schefflera abyssinica</i> , <i>Ficus vasta</i> , <i>Millettia ferruginea</i> and <i>Sesbania sesban</i>	South-western parts of Ethiopia	(Ango et al., 2014; Edo et al., 2017; Gemechu et al., 2021; Hundera et al., 2015; Yakob et al., 2014)
<i>Acacia nilotica</i> , <i>Balanites aegyptiaca</i> , <i>Acacia seyal</i> , <i>Capparis tomentosa</i> , <i>Citrus medica</i> , <i>Carissa edulis</i> , <i>Cordia africana</i> , <i>Ficus sycomorus</i> , <i>Faidherbia albida</i> , <i>Grewia bicolor</i> , <i>Dalbergia melanoxylon</i> <i>Oxytenanthera abyssinica</i> , and <i>Moringa stenopetala</i>	Northern parts of Ethiopia	(Eyasu et al., 2020; Gebrewahid, Teka, et al., 2019; Guyassa et al., 2014)
<i>Acacia abyssinica</i> , <i>Cordia africana</i> , <i>Albizia gummifera</i> , <i>Croton macrostachyus</i> , <i>Faidherbia albida</i> , <i>Erythrina brucei</i> , <i>Rhamnus prinoides</i> <i>Ficus vasta</i> , and <i>Vernonia amygdalina</i>	Central Highlands of Ethiopia	(Duguma & Hager, 2009; Lelamo, 2021; Likassa & Gure, 2014; Yadessa et al., 2009)
<i>Acacia tortilis</i> , <i>Celtis africana</i> , <i>Acacia mellifera</i> , <i>Grewia bicolor</i> , <i>Dichrostachys cinerea</i> <i>Olea europaea</i> , and <i>Balanites aegyptiaca</i>	Mid Rift Valley of Ethiopia	(Shenkute et al., 2012)
<i>Acacia abyssinica</i> , <i>Cordia africana</i> , <i>Albizia gummifera</i> , <i>Erythrina abyssinica</i> and <i>Croton macrostachyus</i>	North-western parts of Ethiopia	(Linger, 2014)

products, which can help families by allowing them to buy food and clothing (Mamo & Asfaw, 2017). Similar to this, native tree species are purposefully preserved and cultivated for their fruit-bearing qualities in the northern part of Ethiopia, notably Tigray (Guyassa et al., 2014). Citrus medica and Annona senegalensis are two important native fruit tree species in Sidama, southern Ethiopia (Adane et al., 2019). In Tigray, women and children gather the fruits of the Cordia africana tree, which is planted in fields or backyards, for sale or personal use (Tewolde Berhan et al., 2013).

Farmers can cover unforeseen expenses with the money they receive from their on-farm trees, especially when there are seasonal droughts and off-seasons (Mesele, 2007). Trees support regional and national economies by generating employment opportunities. Selling the fruits of particular tree species present in landscapes of agriculture in the Tigray area of northern Ethiopia allows children and women to earn money (Guyassa et al., 2014). In southern Ethiopia, native multipurpose fodder trees are producing marketable goods and opening up the potential to raise household incomes (Gina et al., 2014). Some of the benefits, including wood, timber, honey, and medicinal significance, have high commercial worth and increase farmers' ability to generate income (Muleta et al., 2011). Farmers utilize trees like Cordia africana as a security that can generate cash flow and as a risk-avoidance choice during inclement weather seasons (Lelamo, 2021).

3.4.2. The roles of agroforestry trees for climate change mitigation and biodiversity conservation

In contrast to conventional agricultural practices that focus on mono-cropping, agroforestry systems can retain a greater amount of carbon from the atmosphere in both plant components and soil, as noted by Mulhollem (2018). Furthermore, the inclusion of perennial trees in these systems serves to further enhance the sequestration process, as highlighted by Negash et al. (2012). By their above- and below-ground biomass, a variety of multi-purpose trees that are planted and sustained on agricultural land owned by farmers are capable of contributing towards carbon sequestration, as indicated by Gebrewahid, Abrehe, et al. (2019). This is not only relevant in terms of the national carbon budget, but also on a global scale and, thus, is an important avenue to pursue (Zomer et al., 2016). Multipurpose trees are crucial in agroforestry systems because they act as methane sinks at the point where soil and decomposing leaves meet (Gina et al., 2014). Consequently, isolated trees on fields could together significantly increase the climate resilience of a green economy plan (Negash & Starr, 2015). Indigenous

agroforestry systems along Ethiopia's southern rift valley escarpment sequestered an average of 67 Mg ha⁻¹ of total biomass carbon stock, with trees making up 39–93% of that stock (Negash & Starr, 2015). According to the findings of Betemariyam et al. (2020), the implementation of home gardens and coffee-based agroforestry systems nearby can effectively mitigate emissions and enhance carbon sequestration in agricultural landscapes. Such systems can be further integrated into diverse mixed cropping practices on various land types, including but not limited to cropland, pastureland, and rangeland, thereby serving as a potential strategy to counteract the hazardous impacts of climate change while concurrently improving microclimatic conditions (Teketay & Tegineh, 1991).

In traditional Ethiopian agroforestry systems, trees grown on several farmlands combined within a certain area create a more forested environment that promotes environmental preservation. Thus, native woody species are preserved and conserved in a crucial way (Gebrewahid, Abrehe, et al., 2019). According to Kabir and Webb (2008), home gardens function as important wildlife sanctuaries because of their primarily arboreal nature and variety of plant species at various strata. Consequently, they serve as an in-situ technique of conservation for biological variety and plant genetic resources, which includes both flora and fauna (Gemechu et al., 2021; Mulia et al., 2018). A common type of agroforestry, home gardens are a large-scale land use system that can reduce strain on natural forests and preserve biodiversity (Kabir & Webb, 2008; Legesse & Negash, 2021) (Gebrewahid and Abrehe, 2019).

3.4.3. The roles of agroforestry trees for conserving and improving soil fertility

Native agroforestry trees have been recognized in traditional agroforestry systems for their capacity to restore fertility to degraded land, thereby increasing crop yields (Lelamo, 2021). The islands of fertility or centers of variance in soil qualities that trees typically form (Dagar et al., 2020) are partly due to the development of symbiotic relationships with specific soil bacteria, rhizobia, and arbuscular mycorrhizal fungus (Asfaw & Ågren, 2007). These advantages are linked to in-situ activities such as nitrogen cycling, root activity, and litterfall (Yadessa et al., 2009). As Kanshie (2002) noted for trees integrated into Gedeo's agroforestry systems, mulching with tree leaves and short shoots of species like Ficus sur and Cordia africana also contributes to managing soil fertility (Likassa & Gure, 2014), which helps preserve soils and promote organic matter.

The research conducted by Teketay and Tegineh (1991) in Ethiopia has demonstrated the potential of

certain species such as *Cordia africana*, *Millettia ferruginea*, and *Croton macrostachyus* to enhance soil fertility in traditional agroforestry systems. This finding has been supported by various studies conducted by Gindaba et al. (2005), Hailu et al. (2000), Lameso and Bekele (2020), and Mamo and Asfaw (2017). Although *Oxytenanthera abyssinica* and *Dalbergia melanoxylon* are also highly valued on smallholder farms, *Cordia africana* and *Millettia ferruginea* have been recognized as the most effective species for improving soil fertility in North Central and Southern Ethiopia (Gebrewahid, Teka, et al., 2019; Hailu et al., 2000; Kiros et al., 2015).

3.5. Crop rotation and intercropping

In most of Ethiopia, extension programmes routinely include the promotion of crop rotation (Ashango & Mesene, 2019). Several studies have reported (Abera et al., 2009; Alemayehu et al., 2020; Degu et al., 2019; Gorfu et al., 2000) that crop rotation enhances soil structure, reduces soil degradation, and boosts yields. . . Tadesse et al. (2013) found that an increase in soil organic matter enhances nutrition and water retention while lowering the requirement for synthetic fertilisers. Crop rotation effectively addresses climate change mitigation and adaptation (Tilahun et al., 2023). Crop rotation has the potential to enhance nutrient management by decreasing the usage of nitrogen fertiliser and the associated greenhouse gas (GHG) emissions from the production, transportation, and consumption of chemical fertilisers (Feliciano et al., 2022; Meragiaw, 2017; Teklu et al., 2022).

3.6. Small-scale irrigation

Ethiopia has undertaken widespread implementation of small-scale irrigation systems throughout the country, as reported by Amede (2015). Consequently, the area dedicated to irrigated crop production increased to 1,231,660 hectares in 2013, while the coverage of small-scale irrigation infrastructure expanded from 853,000 hectares in 2009 to 2,084,760 hectares in 2013, as documented by MoA (2014). The availability of water and irrigation infrastructure significantly enhances crop and forage yield by elevating biomass, as observed by Van Den Berg and Ruben (2006). In a country that is susceptible to drought and has limited water resources, the provision of irrigation infrastructure and adequate water for agricultural and livestock production can also extend the number of cropping seasons and reduce the risks associated with rain-fed agriculture and livestock rearing, thereby potentially affecting adaptation to climate change and food security, as noted by Bacha

et al. (2011). Effective water management is critical for enhanced agronomic practices and efficient water consumption in irrigated agriculture (Tesfaye & Nayak, 2023). It is therefore imperative to provide appropriate training in agronomy and water management, along with assistance in constructing irrigation infrastructure. In the absence of irrigation, various rainwater harvesting techniques have been employed by many farmers, such as in-field harvesting and more sophisticated methods like roof harvesting, as highlighted by Marie et al. (2020). These methods can aid farmers in conserving water, thereby rendering their crops more resilient to changing weather conditions, particularly mid-season dry spells (Amede, 2015; Bryan et al., 2009).

3.7. Crop diversification and improved variety popularization

The Ethiopian government, along with foreign organisations and non-governmental organisations (NGOs), is working to ensure food security through the development and dissemination of new crops and crop varieties at the community and family level (FAO, 2016). Initiatives to popularise crop varieties are implemented through government and non-governmental organisation programmes as the Eastern Africa Agricultural Productivity Project (EAAPP), SLM, and the Agricultural Growth Programme (AGP) (FAO, 2016). Crop varieties that are resistant to pests and diseases not only lower carbon emissions but also lessen the need for pesticides and the frequency of in-field applications (Warra & Prasad, 2020). In addition to decreasing carbon emissions, pest and disease-resistant crop varieties also lessen the need for pesticides and the frequency of in-field applications (FAO, 2016). Based on the research conducted by Geffersa (2023), it was determined that the use of improved crop varieties, effective inorganic fertilisers, and crop rotation with legumes were important elements in raising agricultural productivity (Figure 3). On the other hand, improved animal husbandry and small-scale irrigation were shown to have moderate weights in relation to the productivity target.

3.8. Traditional climate smart agriculture practices in Ethiopia

Ethiopia has embraced and put into effect a variety of traditional CSA techniques (Tekeste, 2021). These include the Ankober Manure Management (Regassa, 2001), Traditional Agroforestry in Gedeo Zone (Degefa, 2016), East Shewa Zone (Endale et al., 2017), East Wollega Zone (Urgessa Waktola et al., 2021), West Gojam Zone (Chemura et al., 2021), Konso Cultural




Figure 3. Productivity performance of the top five CSA practices. Source: Girma et al, (2023)

Landscape (Gashure & Wana, 2023), Hararghe Highland Traditional Soil and Water Conservation (Bekele & Drake, 2003), Hararghe Cattle Fattening (Dinku, 2019), and Traditional Small-Scale Irrigation (Amede, 2015).

Derashe (Sagandoye Valley) special woreda in SNNP Regional State is one of the locations in Ethiopia where traditional conservation agriculture is practiced (Ashango & Mesene, 2019). Rainfall in the Derashe District's Sagandoye Valley is erratic, with early beginning, dry intervals, and early cessation. So, the problem of food insecurity brought on by poor agricultural productivity and production is a significant one. Due to this difficulty, farmers in the valley have for a long time engaged in conventional conservation agriculture on an estimated 11,000 hectares (Ashango & Mesene, 2019). Sorghum and maize are cultivated using this traditional method without the need to till the ground. Using pointed sticks, seeds are planted in rows. Even in the dry season, when there are no crops on the farm, weeding is routinely done (Muluneh et al., 2022). Crop remnants are put down on the ground in a rectangular pattern after harvesting to retain moisture from the rain (Muluneh et al., 2022). The farm is off-limits to animals, and there is no crop residue removal at all. Crop rotation and intercropping are not, however, practiced consistently, and the extension service must provide support for their promotion (Ashango & Mesene, 2019).

In numerous woredas in the Benishangul-Gumuz (Mosissa & Atinafu, 2021) and Gambella region (Nigussie et al., 2020), smallholder farmers also practice traditional conservation agriculture. The hoe is the primary conventional tool used in this situation to plant

seeds without frequently plowing the ground. Contrary to the Derashe District, traditional conservation agriculture does not include crop rotation in the Benishangul-Gumuz and Gambella regions (Mersha et al., 2022).

In an SNNP Regional State near Derashe Special Woreda, there is a place called the Konso Cultural Landscape (Assoma, 2010). The terrain in the area is steep, and soil erosion is the main cause of environmental damage (Assoma, 2010). Terracing, agroforestry, and manure management are practices used by farmers in Konso to reliably provide strong harvests and preserve the integrity of the soil (Gashure & Wana, 2023). This time-honored method of soil conservation has helped assist in adaptation to climate change while significantly reducing soil erosion (Gashure et al., 2022). The Konso Cultural Landscape is currently included as a UNESCO World Heritage Site as a result of its historical land terracing processes (Gashure et al., 2022).

Farmers in the Hararghe highlands utilize traditional soil and water conservation techniques (Bekele & Drake, 2003; Daba, 2003), similar to those used in the Konso special woreda. Farmers in the Hararghe zones practice small-scale water harvesting and river diversion in addition to soil conservation, which they use to produce irrigated crops (Alemu et al., 2017). Through a cut-and-carry approach, Hararghe also engages in traditional animal fattening (mixed crop-livestock agriculture) (Abebe, 2019; Dinku, 2019; Dinku et al., 2019). Furthermore, traditional agroforestry techniques are still used in regions like East Showa, East and West Wollega, and West Gojam Zone, particularly in the Bure and Wonberima woredas. Farmers in the Amhara Region have

traditionally sprinkled animal dung on crop fields, which has led to notable improvements in crop biomass and output (Lewoyehu & Cardeal, 2021; Tilahun, 2023).

4. Conclusion

Climate Smart Agriculture (CSA) strategies are crucial for enhancing agricultural resilience and food security in Ethiopia, a country vulnerable to climate change impacts. Implementing CSA practices, including climate-resilient crop varieties, conservation agriculture, improved livestock management, and water management systems, can help smallholder farmers adapt to changing conditions, boosting productivity and income. However, challenges such as limited access to finance, technical capacity, and infrastructure hinder their adoption and scaling up. Addressing these requires the active involvement of various stakeholders, including government agencies, international organizations, and private sector actors. The adoption of CSA practices is a critical step towards achieving food security and enhancing agricultural resilience. Long-term planning and sustainability are essential for the success of CSA strategies, and a conducive policy environment supporting CSA practices and stakeholder participation is necessary. Therefore, a conducive policy environment is essential for promoting CSA practices and ensuring the success of CSA strategies in Ethiopia and other vulnerable regions. Efforts to promote CSA must be.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

Abdulbasit Hussein  <http://orcid.org/0000-0002-5067-0104>

Data availability statement

The data was available in the hands of authors confidentially

References

- Abebe, B. (2019). Cattle fattening practices in West Hararghe: Potentials and constraints of beef cattle production in Oromia Regional State, Ethiopia. *Journal of Agricultural Science*, 11(8), 120. <https://doi.org/10.5539/jas.v11n8p120>
- Abebe, T., Wiersum, K. F., & Bongers, F. (2010). Spatial and temporal variation in crop diversity in agroforestry homegardens of southern Ethiopia. *Agroforestry Systems*, 78(3), 309–322. <https://doi.org/10.1007/s10457-009-9246-6>
- Abera, T., Feyisa, D., & Friesen, D. K. (2009). Effects of crop rotation and NP fertilizer rate on grain yield and related characteristics of maize and soil fertility at Bako, Western Oromia, Ethiopia. *East African Journal of Sciences*, 3(1). <https://doi.org/10.4314/eajsci.v3i1.42789>
- Adane, F., Legesse, A., Weldeamanuel, T., & Belay, T. (2019). The contribution of a fruit tree-based agroforestry system for household income to smallholder farmers in Dale District, Sidama Zone, Southern Ethiopia. *Advances in Plants & Agriculture Research*, 9(1), 78–84. <https://doi.org/10.15406/apar.2019.09.00415>
- Adhikari, U., Nejadhashemi, A. P., & Woznicki, S. A. (2015). Climate change and eastern Africa: A review of impact on major crops. *Food and Energy Security*, 4(2), 110–132. <https://doi.org/10.1002/fes3.61>
- Adimassu, Z., Langan, S., & Johnston, R. (2016). Understanding determinants of farmers' investments in sustainable land management practices in Ethiopia: Review and synthesis. *Environment, Development and Sustainability*, 18(4), 1005–1023. <https://doi.org/10.1007/s10668-015-9683-5>
- Agegnehu, G., & Amede, T. (2017). Integrated soil fertility and plant nutrient management in tropical agro-ecosystems: A review. *Pedosphere*, 27(4), 662–680. [https://doi.org/10.1016/S1002-0160\(17\)60382-5](https://doi.org/10.1016/S1002-0160(17)60382-5)
- Agize, M., Demissew, S., & Asfaw, Z. (2013). Indigenous knowledge on management of home gardens and plants in Loma and Gena Bosa districts (weredas) of Dawro Zone, Southern Ethiopia: Plant biodiversity conservation, sustainable utilization and environmental protection. *International Journal of Sciences: Basic and Applied Research (IJSBAR)*, 10(1), 63–99.
- Alemayehu, G., Shibabaw, A., Adgo, E., Asch, F., Freyer, B., & Tejada Moral, M. (2020). Crop rotation and organic matter application restore soil health and productivity of degraded highland crop farms in northwest Ethiopia. *Cogent Food & Agriculture*, 6(1), 1831124. <https://doi.org/10.1080/23311932.2020.1831124>
- Alemu, G., Melka, T., Angasu, B., Degaga, J., & Mechara, E. (2017). Characterization and analysis of farming system in chiro district, West Hararghe zone. *Regional Review Workshop on Completed Research Activities*, 59.
- Allen, J., Pascual, K. S., Romasanta, R. R., Van Trinh, M., Van Thach, T., Van Hung, N., Sander, B. O., & Chivenge, P. (2020). Rice straw management effects on greenhouse gas emissions and mitigation options. *Sustainable Rice Straw Management*, 145–159. <https://doi.org/10.1007/978-3-030-32373-8>
- Altieri, M. A., Nicholls, C. I., Henao, A., & Lana, M. A. (2015). Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*, 35(3), 869–890. <https://doi.org/10.1007/s13593-015-0285-2>
- Amare, D., Wondie, M., Mekuria, W., & Darr, D. (2019). Agroforestry of smallholder farmers in Ethiopia: Practices and benefits. *Small-Scale Forestry*, 18(1), 39–56. <https://doi.org/10.1007/s11842-018-9405-6>

- Amede, T. (2015). Technical and institutional attributes constraining the performance of small-scale irrigation in Ethiopia. *Water Resources and Rural Development*, 6, 78–91. <https://doi.org/10.1016/j.wrr.2014.10.005>
- Andrieu, N., Dumas, P., Hemmerlé, E., Caforio, F., Falconnier, G. N., Blanchard, M., & Vayssières, J. (2021). Ex ante mapping of favorable zones for uptake of climate-smart agricultural practices: A case study in West Africa. *Environmental Development*, 37, 100566. <https://doi.org/10.1016/j.envdev.2020.100566>
- Ango, T. G., Börjeson, L., Senbeta, F., & Hylander, K. (2014). Balancing ecosystem services and disservices: Smallholder farmers' use and management of forest and trees in an agricultural landscape in southwestern Ethiopia. *Ecology and Society*, 19(1). <https://doi.org/10.5751/ES-06279-190130>
- Anshiso, A., Woldeamanuel, T., & Asfaw, Z. (2017). Financial analysis of fruit tree based agroforestry practice in Hadero Tunto Zuria Woreda, Kembata Tembaro Zone, South Ethiopia. *Research Journal of Finance & Accounting*, 8(3), 72–80.
- Araya, T., Cornelis, W. M., Nyssen, J., Govaerts, B., Bauer, H., Gebreegziabher, T., Oicha, T., Raes, D., Sayre, K. D., Haile, M., & Deckers, J. (2011). Effects of conservation agriculture on runoff, soil loss and crop yield under rainfed conditions in Tigray, Northern Ethiopia. *Soil Use and Management*, 27(3), 404–414. <https://doi.org/10.1111/j.1475-2743.2011.00347.x>
- Araya, T., Nyssen, J., Govaerts, B., Deckers, J., & Cornelis, W. M. (2015). Impacts of conservation agriculture-based farming systems on optimizing seasonal rainfall partitioning and productivity on vertisols in the Ethiopian drylands. *Soil and Tillage Research*, 148, 1–13. <https://doi.org/10.1016/j.still.2014.11.009>
- Asfaw, Z., & Ågren, G. I. (2007). Farmers' local knowledge and topsoil properties of agroforestry practices in Sidama, Southern Ethiopia. *Agroforestry Systems*, 71(1), 35–48. <https://doi.org/10.1007/s10457-007-9087-0>
- Ashango, A., & Mesene, M. (2019). Clean Development mechanisms (CDM) and climate smart agriculture (CSA): Role and implication for sustainable natural resource management: Ethiopian context. *Environmental Research*, 11(11), 27–40. <https://doi.org/10.7176/CER/11-11-04>
- Assoma, A. A. (2010). *The "Heritagization" of Konso Cultural Landscape* [Dissertation]. London School of Economics and Political Science.
- Bacha, D., Namara, R., Bogale, A., & Tesfaye, A. (2011). Impact of small-scale irrigation on household poverty: Empirical evidence from the Ambo district in Ethiopia. *Irrigation and Drainage*, 60(1), 1–10. <https://doi.org/10.1002/ird.550>
- Bahru, B. A., & Zeller, M. (2022). Gauging the impact of Ethiopia's productive safety net programme on agriculture: Application of targeted maximum likelihood estimation approach. *Journal of Agricultural Economics*, 73(1), 257–276. <https://doi.org/10.1111/1477-9552.12452>
- Bekele, S. E. (2018). Parkland agroforestry of Ethiopia; key to production, productivity, biodiversity conservation and climate change mitigation: A review. *Open Journal of Forestry*, 8(4), 472. <https://doi.org/10.4236/ojf.2018.84030>
- Bekele, W., & Drake, L. (2003). Soil and water conservation decision behavior of subsistence farmers in the Eastern Highlands of Ethiopia: A case study of the hunde-lafto area. *Ecological Economics*, 46(3), 437–451. [https://doi.org/10.1016/S0921-8009\(03\)00166-6](https://doi.org/10.1016/S0921-8009(03)00166-6)
- Belay, A. D., Kebede, W. M., & Golla, S. Y. (2023). Determinants of climate-smart agricultural practices in smallholder plots: Evidence from Wadla district, northeast Ethiopia. *International Journal of Climate Change Strategies and Management*, 15(5), 619–637. <https://doi.org/10.1108/IJCCSM-06-2022-0071>
- Berhanu, K., & Poulton, C. (2014). The political economy of agricultural extension policy in Ethiopia: Economic growth and political control. *Development Policy Review*, 32(s2), s197–s213. <https://doi.org/10.1111/dpr.12082>
- Betemariyam, M., Negash, M., & Worku, A. (2020). Comparative analysis of carbon stocks in home garden and adjacent coffee based agroforestry systems in Ethiopia. *Small-Scale Forestry*, 19(3), 319–334. <https://doi.org/10.1007/s11842-020-09439-4>
- Birhane, E. (2014). *Agroforestry governance in Ethiopia. Consultancy Report*.
- Brown, B., Nuberg, I., & Llewellyn, R. (2017). Stepwise frameworks for understanding the utilisation of conservation agriculture in Africa. *Agricultural Systems*, 153, 11–22. <https://doi.org/10.1016/j.agry.2017.01.012>
- Bryan, E., Deressa, T. T., Gbetibouo, G. A., & Ringler, C. (2009). Adaptation to climate change in Ethiopia and South Africa: Options and constraints. *Environmental Science & Policy*, 12(4), 413–426. <https://doi.org/10.1016/j.envsci.2008.11.002>
- Burayou, W., Mesfin, T., & Debele, T. (2002). A review of tillage management research on maize in Ethiopia. *Enhancing the contribution of maize to food security in Ethiopia: Proceedings of the second national maize workshop of Ethiopia*, Addis Ababa, Ethiopia (pp. 71–75). EARO/CIMMYT.
- Campbell, B. M. (2017). Climate-smart agriculture- what is it. *Rural 21: The International Journal for Rural Development*, 51(4), 14–16.
- Chemura, A., Yalew, A. W., & Gornott, C. (2021). Quantifying agroforestry yield buffering potential under climate change in the smallholder maize farming systems of Ethiopia. *Frontiers in Agronomy*, 3, 609536. <https://doi.org/10.3389/fagro.2021.609536>
- Conway, D., & Schipper, E. L. F. (2011). Adaptation to climate change in Africa: Challenges and opportunities identified from Ethiopia. *Global Environmental Change*, 21(1), 227–237. <https://doi.org/10.1016/j.gloenvcha.2010.07.013>
- Daba, S. (2003). An investigation of the physical and socio-economic determinants of soil erosion in the Hararghe Highlands, eastern Ethiopia. *Land Degradation & Development*, 14(1), 69–81. <https://doi.org/10.1002/ldr.520>
- Dagar, J. C., Gupta, S. R., & Teketay, D. (2020). *Agroforestry for Degraded Landscapes*. Springer.
- Darcha, G., Birhane, E., & Abadi, N. (2015). Woody species diversity in *Oxytenanthera abyssinica* based homestead agroforestry systems of serako, northern Ethiopia. *Journal of Natural Sciences Research*, 5(9), 18–27.
- Degefa, S. (2016). Home garden agroforestry practices in the Gedeo zone, Ethiopia: A sustainable land management system for socio-ecological benefits. *Socio-Ecological Production Landscapes and Seascapes (SEPLS) in Africa*, 28.
- Degu, M., Melese, A., & Tena, W. (2019). Effects of soil conservation practice and crop rotation on selected soil

- physicochemical properties: The case of Dembecha District, Northwestern Ethiopia. *Applied & Environmental Soil Science*, 2019, 1–14. <https://doi.org/10.1155/2019/6910879>
- Derpsch, R., Friedrich, T., Kassam, A., & Li, H. (2010). Current status of adoption of no-till farming in the world and some of its main benefits. *International Journal of Agricultural and Biological Engineering*, 3(1), 1–25.
- Dinesh, D., Frid-Nielsen, S., Norman, J., Mutamba, M., Loboguerrero Rodriguez, A. M., & Campbell, B. M. (2015). *Is climate-smart agriculture effective? A review of selected cases. CCAFS working paper.*
- Dinku, A. (2019). Assessment of constraints and opportunities in small-scale beef cattle fattening business: Evidence from the West Hararghe Zone of Ethiopia. *International Journal of Veterinary Science and Research*, 5 (2), 58–68. <https://doi.org/10.17352/ijvsr.000042>
- Dinku, A., Abebe, B., Lemma, A., & Shako, M. (2019). Beef cattle value chain analysis: Evidence from West Hararghe Zone of Ethiopia. *International Journal of Agricultural Science and Food Technology*, 5(1), 077–087. <https://doi.org/10.17352/2455-815X.000046>
- Diriba, A., Dekeba, S., & Gizaw, W. (2021). Assessment of existing agroforestry practices in West Hararghe Zone of Oromia region. *International Journal of Research - Granthaalayah*, 9(12), 286–302. <https://doi.org/10.29121/granthaalayah.v9.i12.2021.4436>
- Diro, S., Tesfaye, A., & Erko, B. (2022). Determinants of adoption of climate-smart agricultural technologies and practices in the coffee-based farming system of Ethiopia. *Agriculture & Food Security*, 11(1), 42. <https://doi.org/10.1186/s40066-022-00385-2>
- Djenontin, J. A., Amidou, M., & Baco, N. M. (2004). Diagnostic gestion du troupeau: gestion des ressources pastorales dans les départements de l'Alibori et du Borgou au nord Bénin. *Bulletin de La Recherche Agronomique Du Bénin*, 43(1), 27–31.
- Duguma, L. A., & Hager, H. (2009). Forest products scarcity perception and response by tree planting in the rural landscapes: Farmers, views in central highlands of Ethiopia. *Ekologia*, 28(2), 158–169. https://doi.org/10.4149/ekol_2009_02_158
- Ebabu, M., & Okoyo, N. (2017). Assessment of farmers' climate information need and adoption of climate smart agricultural practices in lasta district, North Wollo Zone, Amhara National Regional State, Ethiopia. Haramaya University.
- Edo, G. Y., Gebremedih, K. G., Woldesenbet, A. F., & Guta, K. K. (2017). Growth performance of some multipurpose tree species around the homesteads in Gimbo District, Southwestern Ethiopia. *Agriculture, Forestry and Fisheries*, 6(1), 1. <https://doi.org/10.11648/j.aff.20170601.11>
- Endale, Y., Derero, A., Argaw, M., & Muthuri, C. (2017). Farmland tree species diversity and spatial distribution pattern in semi-arid East Shewa, Ethiopia. *Forests, Trees and LiveLihoods*, 26(3), 199–214. <https://doi.org/10.1080/14728028.2016.1266971>
- Eyasu, G., Tolera, M., & Negash, M. (2020). Woody species composition, structure, and diversity of homegarden agroforestry systems in southern Tigray, Northern Ethiopia. *Heliyon*, 6(12), e05500. <https://doi.org/10.1016/j.heliyon.2020.e05500>
- Fayera, T. (2022). Climate variability and the responses of crop yields to agricultural drought in the East Wollega Zone, Oromia National Regional State, Ethiopia. *IOP Conference Series: Earth and Environmental Science*, 1016 (1), 12002. <https://doi.org/10.1088/1755-1315/1016/1/012002>
- Feliciano, D., Recha, J., Ambaw, G., MacSween, K., Solomon, D., & Wollenberg, E. (2022). Assessment of agricultural emissions, climate change mitigation and adaptation practices in Ethiopia. *Climate Policy*, 22(4), 427–444. <https://doi.org/10.1080/14693062.2022.2028597>
- Fentie, A., & Beyene, A. D. (2019). Climate-smart agricultural practices and welfare of rural smallholders in Ethiopia: Does planting method matter? *Land Use Policy*, 85, 387–396. <https://doi.org/10.1016/j.landusepol.2019.04.020>
- Gashaw, T. (2015). The implications of watershed management for reversing land degradation in Ethiopia. *Research Journal of Agriculture and Environmental Management*, 4 (1), 5–12.
- Gashure, S., & Wana, D. (2023). Spatiotemporal climate variability and trends in UNESCO designated cultural landscapes of Konso, Ethiopia. *African Geographical Review*, 42(2), 107–124. <https://doi.org/10.1080/19376812.2021.1997611>
- Gashure, S., Wana, D., & Samimi, C. (2022). Impacts of climate variability and climate-smart agricultural practices on crop production in UNESCO designated cultural landscapes of Konso, Ethiopia. *Theoretical and Applied Climatology*, 150(3–4), 1495–1511. <https://doi.org/10.1007/s00704-022-04244-9>
- Gebretsadik, T., & Negash, D. (2016). Assessment of major Honey bee flora resources on selected districts of Sidama and Gedeo zones of South nations nationalities and peoples regional state, Ethiopia. *Journal of Agricultural Economics, Extension and Rural Development*, 4(4), 49–63. <https://doi.org/10.29121/granthaalayah.v4.i4.2016.2754>
- Gebrewahid, Y., Abrehe, S., & Tejada Moral, M. (2019). Biodiversity conservation through indigenous agricultural practices: Woody species composition, density and diversity along an altitudinal gradient of Northern Ethiopia. *Cogent Food & Agriculture*, 5(1), 1700744. <https://doi.org/10.1080/23311932.2019.1700744>
- Gebrewahid, Y., Teka, K., Gebre-Egziabhier, T.-B., Tewolde Berhan, S., Bihane, E., Eyasu, G., & Meresa, E. (2019). Dispersed trees on smallholder farms enhance soil fertility in semi-arid Ethiopia. *Ecological Processes*, 8(1), 1–8. <https://doi.org/10.1186/s13717-019-0190-8>
- Gebru, B. M., Wang, S. W., Kim, S. J., & Lee, W.-K. (2019). Socio-ecological niche and factors affecting agroforestry practice adoption in different agroecologies of southern Tigray, Ethiopia. *Sustainability*, 11(13), 3729. <https://doi.org/10.3390/su11133729>
- Geffersa, A. G. (2023). Does cooperative membership enhance inorganic fertilizer use intensity? Panel data evidence from maize farmers in Ethiopia. *Annals of Public & Cooperative Economics*.
- Gemechu, H. W., Lemessa, D., & Jiru, D. B. (2021). A comparative analysis of indigenous and exotic tree species management practices in agricultural landscapes of Southwest Ethiopia. *Trees, Forests and People*, 4, 100059. <https://doi.org/10.1016/j.tfp.2020.100059>

- Gina, T. G., Nigatu, L., & Animut, G. (2014). Biodiversity of indigenous multipurpose fodder trees of wolayta zone, southern Ethiopia: Ecological and socio-economic importance. *International Journal of Emerging Technology & Advanced Engineering*, 4(5), 494–503.
- Gindaba, J., Rozanov, A., & Negash, L. (2005). Trees on farms and their contribution to soil fertility parameters in Badessa, eastern Ethiopia. *Biology and Fertility of Soils*, 42(1), 66–71. <https://doi.org/10.1007/s00374-005-0859-2>
- Gorfu, A., Girma, K., Tanner, D. G., Taa, A., & Maru, S. (2000). Effect of crop rotation and fertilizer application on wheat yield performance across five years at two locations in southeastern Ethiopia. *Eleventh Regional Wheat Workshop for Eastern, Central and Southern Africa, Addis Ababa, Ethiopia* (pp. 264–274).
- Guyassa, E., Raj, A. J., Gidey, K., & Tadesse, A. (2014). Domestication of indigenous fruit and fodder trees/shrubs in dryland agroforestry and its implication on food security. *International Journal of Ecosystem*, 4(2), 83–88.
- Haile, K. K., Tirivayi, N., & Tesfaye, W. (2019). Farmers' willingness to accept payments for ecosystem services on agricultural land: The case of climate-smart agroforestry in Ethiopia. *Ecosystem Services*, 39, 100964. <https://doi.org/10.1016/j.ecoser.2019.100964>
- Hailu, T., Negash, L., & Olsson, M. (2000). *Milletia ferruginea* from southern Ethiopia: Impacts on soil fertility and growth of maize. *Agroforestry Systems*, 48(1), 9–24. <https://doi.org/10.1023/A:1006274912762>
- Hörner, D., & Wollni, M. (2021). Integrated soil fertility management and household welfare in Ethiopia. *Food Policy*, 100, 102022. <https://doi.org/10.1016/j.foodpol.2020.102022>
- Hulme, M., Doherty, R., Ngara, T., New, M., & Lister, D. (2001). African climate change: 1900–2100. *Climate Research*, 17(2), 145–168. <https://doi.org/10.3354/cr017145>
- Hundera, K., Honnay, O., Aerts, R., & Muys, B. (2015). The potential of small exclosures in assisting regeneration of coffee shade trees in South-Western Ethiopian coffee forests. *African Journal of Ecology*, 53(4), 389–397. <https://doi.org/10.1111/aje.12203>
- Jogdand, O. K. (2020). Study on the effect of global warming and greenhouse gases on environmental system. In *Green chemistry and sustainable technology* (pp. 275–306). Apple Academic Press.
- Kabir, M. E., & Webb, E. L. (2008). Can homegardens conserve biodiversity in Bangladesh? *Biotropica*, 40(1), 95–103. <https://doi.org/10.1111/j.1744-7429.2007.00346.x>
- Kanshie, T. K. (2002). *Five thousand years of sustainability?: A case study on Gedeo land use (Southern Ethiopia)*. Wageningen University and Research.
- Kassam, A., Friedrich, T., Shaxson, F., & Pretty, J. (2009). The spread of conservation agriculture: Justification, sustainability and uptake. *International Journal of Agricultural Sustainability*, 7(4), 292–320. <https://doi.org/10.3763/ijas.2009.0477>
- Ketema, H., & Yimer, F. (2014). Soil property variation under agroforestry based conservation tillage and maize based conventional tillage in Southern Ethiopia. *Soil and Tillage Research*, 141, 25–31. <https://doi.org/10.1016/j.still.2014.03.011>
- Khelifa, R., Mahdjoub, H., Baaloudj, A., Cannings, R. A., & Samways, M. J. (2021). Effects of both climate change and human water demand on a highly threatened damselfly. *Scientific Reports*, 11(1), 7725. <https://doi.org/10.1038/s41598-021-86383-z>
- Kiros, G., Fisseha, I., & Abraham, M. (2015). Evaluation of locally available fertilizer tree or shrub species in Gozamin Woreda, north Central Ethiopia. *Research Journal of Agriculture and Environmental Management*, 4, 164–168.
- Kolberg, D., Persson, T., Mangerud, K., & Riley, H. (2019). Impact of projected climate change on workability, attainable yield, profitability and farm mechanization in Norwegian spring cereals. *Soil and Tillage Research*, 185, 122–138. <https://doi.org/10.1016/j.still.2018.09.002>
- Lameso, L., & Bekele, W. (2020). Farmers local knowledge on niche selection, management strategies and uses of cordia africana tree in agroforestry practices of Sidama zone, southern Ethiopia. *American Journal of Agriculture and Forestry*, 8(6), 258–264. <https://doi.org/10.11648/j.ajaf.20200806.14>
- Legesse, A., & Negash, M. (2021). Species diversity, composition, structure and management in agroforestry systems: The case of Kachabira district, Southern Ethiopia. *Heliyon*, 7(3), e06477. <https://doi.org/10.1016/j.heliyon.2021.e06477>
- Lelamo, L. L. (2021). A review on the indigenous multipurpose agroforestry tree species in Ethiopia: Management, their productive and service roles and constraints. *Heliyon*, 7(9), e07874. <https://doi.org/10.1016/j.heliyon.2021.e07874>
- Lewoyehu, M., & Cardeal, Z. (2021). Evaluation of drinking water quality in rural area of Amhara region, Ethiopia: The case of Mecha district. *Journal of Chemistry*, 2021, 1–11. <https://doi.org/10.1155/2021/9911838>
- Liben, F. M., Tadesse, B., Tola, Y. T., Wortmann, C. S., Kim, H. K., & Mupangwa, W. (2018). Conservation agriculture effects on crop productivity and soil properties in Ethiopia. *Agronomy Journal*, 110(2), 758–767. <https://doi.org/10.2134/agronj2017.07.0384>
- Likassa, E., & Gure, A. (2014). Diversity of shade tree species in smallholder coffee farms of western Oromia, Ethiopia. *African Journal of Geo-Science Research*, 3(1), 1–3.
- Linger, E. (2014). Agro-ecosystem and socio-economic role of homegarden agroforestry in Jabithenan District, North-Western Ethiopia: Implication for climate change adaptation. *Springer Plus*, 3(1), 1–9. <https://doi.org/10.1186/2193-1801-3-154>
- Lipper, L., Thornton, P., Campbell, B. M., Baedeker, T., Braimoh, A., Bwalya, M., Caron, P., Cattaneo, A., Garrity, D., Henry, K., Hottle, R., Jackson, L., Jarvis, A., Kossam, F., Mann, W., McCarthy, N., Meybeck, A., Neufeldt, H., & Tibu, A. (2014). Climate-smart agriculture for food security. *Nature Climate Change*, 4(12), 1068–1072. <https://doi.org/10.1038/nclimate2437>
- Mamo, D., & Asfaw, Z. (2017). Assessment of farmers' management activities on scattered trees on crop fields at Gemechis district, West Hararge Zone, Oromia, Ethiopia. *International Journal of Agriculture*, 2(1), 41–57.
- Marie, M., Yirga, F., Haile, M., & Tquabo, F. (2020). Farmers' choices and factors affecting adoption of climate change adaptation strategies: Evidence from northwestern Ethiopia. *Heliyon*, 6(4), e03867. <https://doi.org/10.1016/j.heliyon.2020.e03867>

- McMichael, A. J. (2013). Globalization, climate change, and human health. *New England Journal of Medicine*, 368(14), 1335–1343. <https://doi.org/10.1056/NEJMra1109341>
- Mengistie, B. T., Mol, A. P. J., & Oosterveer, P. (2017). Pesticide use practices among smallholder vegetable farmers in Ethiopian Central Rift Valley. *Environment, Development and Sustainability*, 19(1), 301–324. <https://doi.org/10.1007/s10668-015-9728-9>
- Meragiaw, M. (2017). Role of agroforestry and plantation on climate change mitigation and carbon sequestration in Ethiopia. *Journal of Tree Sciences*, 36(1), 1–15. <https://doi.org/10.5958/2455-7129.2017.00001.2>
- Merga, G. (2020). The role of smallscale irrigation schemes in poverty reduction: the case of Sibire Woreda, Western Ethiopia. *Editorial Board*, 9(11), 111.
- Mersha, F., Haji, J., Emana, B., & Mehare, A. (2022). Farm households choices of adaptation strategies to climate variability challenges in Benishangul Gumuz Regional State, Western Ethiopia. *Sustainable Agriculture Research*, 11(4), 50–54. <https://doi.org/10.5539/sar.v11n4p50>
- Mesele, N. (2007). Tree's management and livelihoods in Gedeo's Agroforests, Ethiopia. *Forest. Trees and Livelihood*, 17(2), 158–167. <https://doi.org/10.1080/14728028.2007.9752591>
- Mhired, D. A., Dagnew, D. C., Guzman, C. D., Alemie, T. C., Zegeye, A. D., Tebebu, T. Y., Langendoen, E. J., Zaitchik, B. F., Tilahun, S. A., & Steenhuis, T. S. (2020). A nine-year study on the benefits and risks of soil and water conservation practices in the humid highlands of Ethiopia: The Debre Mawi watershed. *Journal of Environmental Management*, 270, 110885. <https://doi.org/10.1016/j.jenvman.2020.110885>
- Mohamed-Brahmi, A., Tsiokos, D., Ben Saïd, S., Boudalia, S., Smeti, S., Bousbia, A., Gueroui, Y., Boudebbouz, A., Anastasiadou, M., & Symeon, G. K. (2022). Challenges and opportunities of the Mediterranean indigenous bovine populations: Analysis of the different production systems in Algeria, Greece, and Tunisia. *Sustainability*, 14(6), 3356.
- Molyneux, N., Da Cruz, G. R., Williams, R. L., Andersen, R., & Turner, N. C. (2012). Climate change and population growth in Timor Leste: Implications for food security. *AMBIO: A Journal of the Human Environment*, 41(8), 823–840. <https://doi.org/10.1007/s13280-012-0287-0>
- Mosissa, D., & Atinafu, H. (2021). Applied ethnobotany: People, medicinal plants use and conservation practices in Benishangul Gumuz Regional State of Ethiopia: The future cursed natural resource in the Region. *International Journal of Pharmaceutical Sciences*, 1(1), 18–30.
- Muleta, D., Assefa, F., Nemomissa, S., & Granhall, U. (2011). Socioeconomic benefits of shade trees in coffee production systems in Bonga and Yayuhurumu districts, southwestern Ethiopia: Farmers' perceptions. *Ethiopian Journal of Education and Sciences*, 7(1), 39–55.
- Mulhollem, J. (2018). Agroforestry systems may play vital role in mitigating climate change. *Penn State News*, February, 1.
- Mulia, R., Simelton, E., Le, T. T., Pham, T. V., & Do, T. H. (2018). Native and endangered timber tree species in home gardens of northeast and North Central Vietnam. *Biodiversity International Journal*, 2(2), 40–43. <https://doi.org/10.15406/bij.2018.02.00041>
- Muluneh, S. A., Tekalign, S., & Asefaw, S. (2022). *Practices of climate smart agriculture and impacts on the household food security in rural kebeles of Dire Dawa administration, Ethiopia*. Haramaya University.
- Mutenje, M. J., Farnworth, C. R., Stirling, C., Thierfelder, C., Mupangwa, W., & Nyagumbo, I. (2019). A cost-benefit analysis of climate-smart agriculture options in Southern Africa: Balancing gender and technology. *Ecological Economics*, 163, 126–137. <https://doi.org/10.1016/j.ecolecon.2019.05.013>
- Negash, M. (2007). Trees management and livelihoods in Gedeo's agroforests, Ethiopia. *Forests, Trees and Livelihoods*, 17(2), 157–168. <https://doi.org/10.1080/14728028.2007.9752591>
- Negash, M., & Starr, M. (2015). Biomass and soil carbon stocks of indigenous agroforestry systems on the south-eastern Rift Valley escarpment, Ethiopia. *Plant and Soil*, 393(1), 95–107. <https://doi.org/10.1007/s11104-015-2469-6>
- Negash, M., Yirdaw, E., & Luukkanen, O. (2012). Potential of indigenous multistrata agroforests for maintaining native floristic diversity in the south-eastern Rift Valley escarpment, Ethiopia. *Agroforestry Systems*, 85(1), 9–28. <https://doi.org/10.1007/s10457-011-9408-1>
- Ngosong, C., Okolle, J. N., & Tening, A. S. (2019). Mulching: A sustainable option to improve soil health. *Soil Fertility Management for Sustainable Development*, 231–249.
- Nigussie, E., Betebo, T., & Yousuf, J. (2020). The role of indigenous knowledge in conservation agriculture: The case of lare district in Gambella Region, Ethiopia. *Innovative Systems Design and Engineering*, 11(1), 8–18.
- Ntawuruhunga, D., Ngowi, E. E., Mangi, H. O., Salanga, R. J., & Shikuku, K. M. (2023). Climate-smart agroforestry systems and practices: A systematic review of what works, what doesn't work, and why. *Forest Policy and Economics*, 150, 102937. <https://doi.org/10.1016/j.forpol.2023.102937>
- Nyasimi, M., Amwata, D., Hove, L., Kinyangi, J., & Wamukoya, G. (2014). *Evidence of impact: climate-smart agriculture in Africa*. CCAFS working paper.
- Paul, C. J., & Weinthal, E. (2019). The development of Ethiopia's climate resilient green economy 2011–2014: Implications for rural adaptation. *Climate and Development*, 11(3), 193–202. <https://doi.org/10.1080/17565529.2018.1442802>
- Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., & Cassman, K. G. (2014). Limited potential of no-till agriculture for climate change mitigation. *Nature Climate Change*, 4(8), 678–683. <https://doi.org/10.1038/nclimate2292>
- Regassa, S. (2001). Decision making on manure use and fallowing as soil fertility maintenance techniques in the northern highlands of Ethiopia: The case of Ankober District.
- Shenkute, B., Hassen, A., Assafa, T., Amen, N., & Ebro, A. (2012). Identification and nutritive value of potential fodder trees and shrubs in the mid Rift Valley of Ethiopia.
- Shonde, Y. (2017). Livelihood contributions of Moringa tree based agroforestry practices in Konso district, southern Ethiopia. *Journal of Resources Development and Management*, 36, 2422–8397.
- Srivastava, R. K., & Srivastava, R. K. (2020). Urban agglomerates under climate change induced risk. *Managing Urbanization, Climate Change and Disasters in South Asia*, 199–250.

- Tadesse, T., Dechassa, N., Bayu, W., & Gebeyehu, S. (2013). Effects of farmyard manure and inorganic fertilizer application on soil physico-chemical properties and nutrient balance in rain-fed lowland rice ecosystem. *American Journal of Plant Sciences*, 04(2), 309–316. <https://doi.org/10.4236/ajps.2013.42041>
- Taylor, M. (2018). Climate-smart agriculture: What is it good for? *The Journal of Peasant Studies*, 45(1), 89–107. <https://doi.org/10.1080/03066150.2017.1312355>
- Tekeste, K. (2021). Climate-smart agricultural (CSA) practices and its implications to food security in Siyadebrina Wayu District, Ethiopia. *African Journal of Agricultural Research*, 17(1), 92–103. <https://doi.org/10.5897/AJAR2020.15100>
- Teketay, D., & Tegineh, A. (1991). Shade trees of coffee in Harerge, Eastern Ethiopia. *International Tree Crops Journal*, 7(1–2), 17–27. <https://doi.org/10.1080/01435698.1991.9752899>
- Teklu, A., Simane, B., & Bezabih, M. (2022). Effectiveness of climate-smart agriculture innovations in smallholder agriculture system in Ethiopia. *Sustainability*, 14(23), 16143. <https://doi.org/10.3390/su142316143>
- Tesfaye, G., Alamirew, T., Kebede, A., & Zeleke, G. (2018). Institutional functionality in participatory integrated watershed development of Tana sub-basin, Ethiopia. *The Land*, 7(4), 130. <https://doi.org/10.3390/land7040130>
- Tesfaye, T., & Nayak, D. (2023). Climate change adaptation measures by farm households in Gedeo zone, Ethiopia: An application of multivariate analysis approach. *Environment, Development and Sustainability*, 25(4), 3183–3209. <https://doi.org/10.1007/s10668-022-02185-x>
- Tewelde Berhan, S., Remberg, S. F., Abegaz, K., Narvhus, J., Abay, F., & Wicklund, T. (2013). Ferric reducing antioxidant power and total phenols in Cordia africana fruit. *African Journal of Biochemistry Research*, 7(11), 215–224. <https://doi.org/10.5897/AJBR2013.0692>
- Tilahun, H. (2023). Assessment on rearing and husbandry practices of indigenous goats in North Shewa Zone, Amhara region, Ethiopia. *Journal of Applied Animal Research*, 51(1), 242–255. <https://doi.org/10.1080/09712119.2023.2185625>
- Tilahun, G., Bantider, A., & Yayeh, D. (2023). Synergies and trade-offs of climate-smart agriculture (CSA) practices selected by smallholder farmers in geshy watershed, Southwest Ethiopia. *Regional Sustainability*, 4(2), 129–138. <https://doi.org/10.1016/j.regsus.2023.04.001>
- Tilahun, S. A., Guzman, C. D., Zegeye, A. D., Ayana, E. K., Collick, A. S., Yitaferu, B., & Steenhuis, T. S. (2014). Spatial and temporal patterns of soil erosion in the semi-humid Ethiopian highlands: A case study of Debre Mawi watershed. *Nile River Basin: Ecohydrological Challenges, Climate Change and Hydropolitics*, 149–163.
- Tsegayea, W., Aredob, D., La Rovere, R., Mwangi, W., Mwabu, G., & Kassie, G. T. (2010). *Adoption and Impact of Conservation Agriculture in Central Ethiopia: Application of Instrumental and Control Function Estimations*.
- Tsegaye, W., Aredo, D., La Rovere, R., Mwangi, W., Kassie, G. T., & Mwabu, G. (2015). *Adoption and impact of conservation agriculture in central Ethiopia*. CIMMYT.
- Urgessa Waktola, T., Fekadu, K., & Okatan, V. (2021). Adoption of coffee shade agroforestry technology and shade tree management in gobu seyo district, east wollega, oromia. *Advances in Agriculture*, 2021, 1–13. <https://doi.org/10.1155/2021/8574214>
- Van Den Berg, M., & Ruben, R. (2006). Small-scale irrigation and income distribution in Ethiopia. *The Journal of Development Studies*, 42(5), 868–880. <https://doi.org/10.1080/00220380600742142>
- Warra, A. A., & Prasad, M. N. V. (2020). African perspective of chemical usage in agriculture and horticulture—their impact on human health and environment. In *Agrochemicals detection, treatment and remediation* (pp. 401–436). Elsevier.
- Wassie, A., & Pauline, N. (2018). Evaluating smallholder farmers' preferences for climate smart agricultural practices in Tehuledere District, northeastern Ethiopia. *Singapore Journal of Tropical Geography*, 39(2), 300–316. <https://doi.org/10.1111/sjtg.12240>
- Wekesa, B. M., Ayuya, O. I., & Lagat, J. K. (2018). Effect of climate-smart agricultural practices on household food security in smallholder production systems: Micro-level evidence from Kenya. *Agriculture & Food Security*, 7(1), 1–14. <https://doi.org/10.1186/s40066-018-0230-0>
- Were, K., Gelaw, A. M., & Singh, B. R. (2016). Smart strategies for enhanced agricultural resilience and food security under a changing climate in sub-Saharan Africa. *Climate Change and Multi-Dimensional Sustainability in African Agriculture: Climate Change and Sustainability in Agriculture*, 431–453.
- Westermann, O., Förch, W., Thornton, P., Körner, J., Cramer, L., & Campbell, B. (2018). Scaling up agricultural interventions: Case studies of climate-smart agriculture. *Agricultural Systems*, 165, 283–293. <https://doi.org/10.1016/j.agsy.2018.07.007>
- Williams, T. O., Mul, M. L., Cofie, O. O., Kinyangi, J., Zougmore, R. B., Wamukoya, G., Nyasimi, M., Mapfumo, P., Speranza, C. I., & Amwata, D. (2015). Climate smart agriculture in the African context.
- Yadessa, A., Itanna, F., & Olsson, M. (2009). Scattered trees as modifiers of agricultural landscapes: The role of waddeessa (*Cordia africana* Lam.) trees in Bako area, Oromia, Ethiopia. *African Journal of Ecology*, 47(s1), 78–83. <https://doi.org/10.1111/j.1365-2028.2008.01053.x>
- Yakob, G., Asfaw, Z., & Zewdie, S. (2014). Wood production and management of woody species in homegardens agroforestry: The case of smallholder farmers in Gimbo district, south west Ethiopia. *International Journal of Natural Sciences Research*, 2(10), 165–175.
- Yebo, B. (2015). Integrated soil fertility management for better crop production in Ethiopia. *International Journal of Soil Science*, 10(1), 1–16. <https://doi.org/10.3923/ijss.2015.1.16>
- Zegeye, H. (2018). Climate change in Ethiopia: Impacts, mitigation and adaptation. *International Journal of Research in Environmental Studies*, 5(1), 18–35.
- Zegeye, A. D., Langendoen, E. J., Stoof, C. R., Tilahun, S. A., Dagnew, D. C., Zimale, F. A., Guzman, C. D., Yitaferu, B., & Steenhuis, T. S. (2016). Morphological dynamics of gully systems in the subhumid Ethiopian Highlands: The Debre Mawi watershed. *Soil*, 2(3), 443–458. <https://doi.org/10.5194/soil-2-443-2016>

- Zerssa, G., Feyssa, D., Kim, D.-G., & Eichler-Löbermann, B. (2021). Challenges of smallholder farming in Ethiopia and opportunities by adopting climate-smart agriculture. *Agriculture*, 11(3), 192. <https://doi.org/10.3390/agriculture11030192>
- Zomer, R. J., Neufeldt, H., Xu, J., Ahrends, A., Bossio, D., Trabucco, A., Van Noordwijk, M., & Wang, M. (2016). Global tree cover and biomass carbon on agricultural land: The contribution of agroforestry to global and national carbon budgets. *Scientific Reports*, 6(1), 29987. <https://doi.org/10.1038/srep29987>
- Zougmore, R., Partey, S., Ouédraogo, M., Omitoyin, B., Thomas, T., Ayantunde, A., Ericksen, P., Said, M., & Jalloh, A. (2016). Toward climate-smart agriculture in West Africa: A review of climate change impacts, adaptation strategies and policy developments for the livestock, fishery and crop production sectors. *Agriculture & Food Security*, 5(1), 1–16. <https://doi.org/10.1186/s40066-016-0075-3>